

Assignment 16

SEPARABILITY, COMPUTATION OF SOME GALOIS GROUPS

1. Let $f \in k[X]$ and let $E \supset k$ be a splitting field of f . We want to prove that f has no multiple root in E if and only if $\gcd_{k[X]}(f, f') = 1$.
 - (a) Let F/k be a field extension and $f, g \in k[X]$. Prove that $\gcd_{k[X]}(f, g) = 1$ if and only if $\gcd_{F[X]}(f, g) = 1$.
 - (b) Write $f = \prod_{i=1}^n (X - \alpha_i)$ in $E[X]$. Establish the formula

$$\prod_{i=1}^n f'(\alpha_i) = \pm \left(\prod_{i < j} (\alpha_i - \alpha_j) \right)^2.$$

- (c) Use the above steps in order to conclude.
2. Let p be a prime number. Consider the polynomial $\varphi_p := \frac{X^p-1}{X-1} \in \mathbb{Q}[X]$ and let $\zeta := e^{\frac{2\pi i}{p}}$. Let $E := \text{Sf}(\varphi_p)$.
 - (a) Prove that φ_p is irreducible in $\mathbb{Q}[X]$ and deduce that $\varphi_p = \text{irr}(\zeta; \mathbb{Q})$.
 - (b) Show that $E = \mathbb{Q}(\zeta)$.
 - (c) Prove that $\text{Gal}(E/\mathbb{Q}) = (\mathbb{Z}/p\mathbb{Z})^\times$.
 3. Let $E = \mathbb{Q}(\sqrt{2}, \sqrt{3})$.
 - (a) Prove that $[E : \mathbb{Q}] = 4$.
 - (b) Show that $\text{Gal}(E/\mathbb{Q}) = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$.
 4. Show that the Galois group of $X^3 - 2 \in \mathbb{Q}[X]$ is isomorphic to S_3 . [*Hint:* Let $E = \text{Sf}(X^3 - 2)$. Find the roots of $X^3 - 2$ in \mathbb{C} . Consider the intermediate extension $\mathbb{Q}(\exp(2\pi i/3))/\mathbb{Q}$ of E and show that $[E : \mathbb{Q}] > 3$.]