

Answer sheet compilation instructions

- Use only black or blue pen.

For open answers:

- Write clearly only **inside** boxes, away from the borders.
- Write a **single** character (number or letter) per box.
- Start writing from the left, leaving empty boxes on the right.

For multiple choice and true/false questions:

- **Fill** the circle for the answer you consider correct (only one answer is correct).
- Remarks and computations have **no** influence on points awarded.
- Any unclear or double marking will be considered as an answer not given (0 points).
- Wrong answers give **negative** points.

Exam instructions

- Turn off your devices and leave them in your bag.
- Only pens and Legi should be on the table.
- Fill last name and Legi number on the answer sheet.
- **Turn this sheet only when instructed to do so.**
- At the end of the exam, take everything except the single answer sheet which you want to submit.



Questions

NumCSE midterm, HS 2017

1. *Cancellation error* [12 P.]

Which of the following expressions can be affected by cancellation errors for some choice of x in the specified interval? (True = affected by cancellation, False = **not** affected by cancellation).

(a) $y = \log_2(x - \sqrt{x-1})$, for $x \in [1, \infty)$. F

(b) $y = \exp(x) - \exp(2(x-1))$, for $x \in (1, 2]$. T

(c) $y = \frac{1 - \cos(x)}{x^2}$, for $x \in (0, \frac{\pi}{2})$. T

(d) $y = x^2 - \sqrt{x^2 + 2}$, for $x \in [2, \infty)$. F

2. Numerical Stability [8 P.]

Let $\tilde{F} : X \mapsto Y$ be an algorithm for the problem $F : X \mapsto Y$.

(a) True or false: backward stability of \tilde{F} implies mixed stability. *True*

(b) True or false: if \tilde{F} is backward stable then, for any $x \in X$, $\tilde{F}(x)$ will be close to $F(x)$. *False*

(c) Among the following, choose the best definition of condition number of F in x :

(i) $\inf_{\Delta x \text{ small}} \left(\frac{\|F(x + \Delta x) - F(x)\|}{\|F(x)\|} \cdot \frac{\|\Delta x\|}{\|x\|} \right)$

(ii) $\sup_{\Delta x \text{ small}} \left(\frac{\|F(x + \Delta x) - F(x)\|}{\|F(x)\|} \cdot \frac{\|\Delta x\|}{\|x\|} \right)$

(iii) $\inf_{\Delta x \text{ small}} \left(\frac{\|F(x + \Delta x) - F(x)\|}{\|F(x)\|} \cdot \frac{\|x\|}{\|\Delta x\|} \right)$

~~(iv)~~ $\sup_{\Delta x \text{ small}} \left(\frac{\|F(x + \Delta x) - F(x)\|}{\|F(x)\|} \cdot \frac{\|x\|}{\|\Delta x\|} \right)$

3. Linear system solution through LU decomposition [9 P.]

Consider the following C++/Eigen function:

```
1  int solve_triang(int n) {
2      using namespace Eigen;
3      MatrixXd A = MatrixXd::Zero(n,n);
4
5      for (int i=0; i < n; i++) {
6          for (int j=i; j < n; j++) {
7              // fill upper triangular part of A
8              A(i,j) = 4;
9          }
10     }
11
12     // fill b
13     VectorXd b = 36 * VectorXd::Ones(n);
14
15     // compute solution to Ax = b
16     VectorXd x = A.fullPivLu().solve(b);
17
18     return x(n-1);
19 }
```

- (a) What is the value returned by `solve_triang(2017)`? 9
- (b) Choose the lowest correct asymptotic complexity of the function as $n \rightarrow \infty$:
- ~~(i)~~ $O(n^3)$
 - (ii) $O(n^2 \log n)$
 - (iii) $O(n^2)$
 - (iv) $O(n \log n)$
- (c) Suppose we replace line 16 with
- ```
VectorXd x = A.triangularView<Upper>().solve(b);
```
- Choose the lowest correct asymptotic complexity of the modified function as  $n \rightarrow \infty$ :
- (i)  $O(n^3)$
  - (ii)  $O(n^2 \log n)$
  - ~~(iii)~~  $O(n^2)$
  - (iv)  $O(n \log n)$

4. Linear Least Squares [12 P.]

Consider the linear least squares problem:

$$\mathbf{x}^* = \arg \min_{\mathbf{x} \in \mathbb{R}^n} \|\mathbf{Ax} - \mathbf{b}\|_2 \quad (1)$$

where  $\mathbf{A} \in \mathbb{R}^{m,n}$  is a large sparse matrix and  $\mathbf{b} \in \mathbb{R}^m$ , for  $m \geq n$  and  $m, n \in \mathbb{N}$ .

(a) State whether the following relations are true or false:

(i)  $\mathcal{N}(\mathbf{A}^\top \mathbf{A}) = \mathcal{N}(\mathbf{A})^\perp$  F

(ii)  $\mathcal{R}(\mathbf{A}^\top \mathbf{A}) = \mathcal{R}(\mathbf{A}^\top)$  T

(iii)  $\mathcal{R}(\mathbf{A}^\top \mathbf{A}) = \mathcal{N}(\mathbf{A})^\perp$  T

(iv)  $\mathcal{N}(\mathbf{A}^\top \mathbf{A}) = \mathcal{R}(\mathbf{A}^\top)^\perp$  T

Here  $\mathcal{R}$  stands for range and  $\mathcal{N}$  for null-space.

(b) If  $\mathbf{x}^*$  is unique, then what is the rank of  $\mathbf{A}$ ? w

(c) Assuming  $\mathbf{A}$  is well-conditioned and a unique solution exists, choose the most efficient method to solve (1) among the following:

- ~~(i)~~ Extended normal equations
- (ii) Singular value decomposition
- (iii) Householder QR decomposition
- (iv) Normal equations

5. *Singular Value Decomposition* [9 P.]

Consider the matrix

$$\mathbf{A} = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 3 & 0 \\ 0 & 0 & 36 & 0 & 0 \\ 0 & 12 & 0 & 0 & 0 \\ 2 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

Let  $\mathbf{U}$ ,  $\mathbf{\Sigma}$ ,  $\mathbf{V}$  be the matrices involved in the **thin** singular value decomposition of  $\mathbf{A}$  (in particular  $\mathbf{A} = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T$ ).

- (a) What is the condition number of  $\mathbf{\Sigma}$ ?  $36/2 = 18$  or  $\infty$
- (b) What is the condition number of  $\mathbf{\Sigma}^T\mathbf{\Sigma}$ ?  $(18)^2 = 324$  or  $\infty$

- (c) What are the dimensions of  $\mathbf{U}$  (number of rows  $\times$  number of columns)?

$$5 \times 4 \text{ or } 5 \times 5$$