

Analysis III

Solutions Serie 13

1. Find the solution of the 2-dimensional Laplace equation (steady heat equation)

$$\nabla^2 u = 0$$

on the disk $D = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 4\}$ with prescribed temperature on the external circumference given by

$$u(x, y) = x^3, \quad (x, y) \in \partial D \left(= \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 4\} \right).$$

[You can use the trigonometric formula

$$\cos^3(\theta) = \frac{3}{4} \cos(\theta) + \frac{1}{4} \cos(3\theta)]$$

Do as follows:

- a) Write the boundary condition in polar coordinates.

Solution:

The polar coordinates are linked to the standard cartesian ones by

$$\begin{cases} x = r \cos(\theta) \\ y = r \sin(\theta) \end{cases}$$

At the boundary the radius is $r = 2$ and the function becomes

$$x^3 = (2 \cos(\theta))^3 = 8 \cos^3(\theta).$$

- b) Solve the problem in polar coordinates, using the methods/formulas learnt in the Lecture notes.

Solution:

The Poisson integral formula provides always the solution, but it is very difficult to get an explicit solution out of it (solving the integral), so we will try to avoid it whenever possible.

In fact, when the boundary condition is easy, it will be also easy to match the coefficients in the Fourier series.

Namely, from the Lecture notes we have a general solution of the Laplace equation in polar coordinates

$$u(r, \theta) = \sum_{n=0}^{+\infty} r^n (A_n \cos(n\theta) + B_n \sin(n\theta)).$$

We impose that on the boundary $r = 2$:

$$u(2, \theta) = \sum_{n=0}^{+\infty} 2^n (A_n \cos(n\theta) + B_n \sin(n\theta)) = 8 \cos^3(\theta).$$

Which amounts in finding the Fourier series of $8 \cos^3(\theta)$:

$$8 \cos^3(\theta) = a_0 + \sum_{n=1}^{+\infty} a_n \cos(n\theta) + \sum_{n=1}^{+\infty} b_n \sin(n\theta)$$

The formula given in the text of the Exercise allows us to avoid computing any integral to get

$$8 \cos^3(\theta) = 6 \cos(\theta) + 2 \cos(3\theta).$$

In terms of the coefficients we had to find this means

$$\begin{cases} B_n = 0 & \forall n \geq 0 \\ A_n = 0 & \forall n \geq 0, n \neq 1, 3 \\ A_1 = \frac{a_1}{2} = 3 \\ A_3 = \frac{a_3}{2^3} = \frac{1}{4} \end{cases}$$

and the solution is

$$u(r, \theta) = rA_1 \cos(\theta) + r^3A_3 \cos(3\theta) = 3r \cos(\theta) + \frac{r^3}{4} \cos(3\theta).$$

c) Express the solution in the standard cartesian coordinates:

$$u(x, y) = ?$$

If we want to express some function in terms of the standard cartesian coordinates (x, y) we are ok with terms like $r, \cos^k(\theta), \sin^k(\theta)$. So in this case we just need to write the function $\cos(3\theta)$ in these terms:

$$\cos(3\theta) = 4 \cos^3(\theta) - 3 \cos(\theta)$$

and

$$\begin{aligned} 3r \cos(\theta) + \frac{r^3}{4} \cos(3\theta) &= 3r \cos(\theta) + \frac{r^3}{4} (4 \cos^3(\theta) - 3 \cos(\theta)) = \\ &= 3(r \cos(\theta)) + (r \cos(\theta))^3 - \frac{3}{4} r^2 (r \cos(\theta)) = 3x + x^3 - \frac{3}{4} (x^2 + y^2)x = \\ &= 3x + x^3 - \frac{3}{4} x^3 - \frac{3}{4} xy^2 = 3x + \frac{1}{4} x^3 - \frac{3}{4} xy^2. \end{aligned}$$

2. Let $R > 0$.

- a) Find the solution $u(r, \theta)$ of the Dirichlet problem on the disk of radius R in polar coordinates:

$$\begin{cases} \nabla^2 u = 0, & 0 \leq r \leq R, 0 \leq \theta \leq 2\pi \\ u(R, \theta) = \sin^2(\theta), & 0 \leq \theta \leq 2\pi. \end{cases}$$

[You can use the trigonometric formula

$$\sin^2(\theta) = \frac{1}{2} - \frac{1}{2} \cos(2\theta)]$$

Solution:

The solution will be

$$u(r, \theta) = \sum_{n=0}^{+\infty} r^n (A_n \cos(n\theta) + B_n \sin(n\theta)),$$

with coefficients found imposing

$$u(R, \theta) = \sum_{n=0}^{+\infty} R^n (A_n \cos(n\theta) + B_n \sin(n\theta)) = \sin^2(\theta).$$

Using the trigonometric formula

$$\sin^2(\theta) = \frac{1}{2} - \frac{1}{2} \cos(2\theta)$$

we obtain coefficients

$$\begin{cases} B_n = 0, & \forall n \geq 0 \\ A_n = 0, & \forall n \geq 0, n \neq 0, 2 \\ A_0 = \frac{1}{2}, \\ A_2 = -\frac{1}{2R^2}. \end{cases}$$

Finally

$$u(r, \theta) = \frac{1}{2} - \frac{r^2}{2R^2} \cos(2\theta).$$

- b) Find the maximum of $u(r, \theta)$. In which point(s) is it reached?

Solution:

The constant $1/2$ plays no role in finding where is the maximum. The other term is a product of functions dependent from different variables. We then just need to maximize these functions independently:

$$\begin{cases} -\cos(2\theta) = 1 \\ r = R \end{cases} \Leftrightarrow \begin{cases} \theta = \frac{\pi}{2}, \frac{3\pi}{2} \\ r = R \end{cases} \rightsquigarrow (r, \theta) = \left(R, \frac{\pi}{2}\right), \left(R, \frac{3\pi}{2}\right) =: P_1, P_2$$

So the maximum is reached in the two points P_1, P_2 and it's equal to

$$u(P_1) = u(P_2) = \frac{1}{2} + \frac{R^2}{2R^2} = \frac{1}{2} + \frac{1}{2} = 1.$$

Remark: the maximum is reached only the boundary. Inside the disk the function is always strictly smaller than this value. You will see in the next lecture that this is not a coincidence.

- c) Express the solution in the standard cartesian coordinates.

Solution:

We transform the term

$$\cos(2\theta) = \cos^2(\theta) - \sin^2(\theta) \rightsquigarrow r^2 \cos(2\theta) = x^2 - y^2$$

and the solution will be

$$u(x, y) = \frac{1}{2} - \frac{1}{2R^2}(x^2 - y^2).$$

3. Let u be the unique harmonic function ($\nabla^2 u = 0$) on the unit disk which on the boundary is

$$u(x, y) = xy + 3.$$

Without computing any integral or using any formula from the script,
answer the following questions:

- a) Find the solution in the standard coordinates $u(x, y)$.

Solution:

The function on the boundary is well defined everywhere in the plane (x, y) . Also, being linear in each variable, it's harmonic:

$$\nabla^2(xy + 3) = \partial_{xx}(xy + 3) + \partial_{yy}(xy + 3) = 0.$$

By uniqueness of the solution for the Dirichlet problem it must be itself the solution:

$$u(x, y) = xy + 3.$$

- b) What's the value in the center of the disk $u(0, 0) = ?$

Solution:

$$u(0, 0) = 0 \cdot 0 + 3 = 3.$$

c) What's the maximum of u and in which point(s) is it reached?

Solution:

To find the maximum is more convenient to write the solution in polar coordinates in this case

$$u(r, \theta) = r^2 \cos(\theta) \sin(\theta) + 3 = \frac{r^2}{2} \sin(2\theta) + 3.$$

Like we did before we need to impose

$$\begin{cases} \sin(2\theta) = 1 \\ r = R = 1 \end{cases} \Leftrightarrow \begin{cases} \theta = \frac{\pi}{4}, \frac{5}{4}\pi \\ r = 1 \end{cases} \rightsquigarrow (r, \theta) = \left(1, \frac{\pi}{4}\right), \left(1, \frac{5}{4}\pi\right) =: Q_1, Q_2.$$

So the maximum is reached in the two points Q_1, Q_2 with value

$$u(Q_1) = u(Q_2) = \frac{1^2}{2} + 3 = \frac{1}{2} + 3 = \frac{7}{2}.$$

4. Prove, without computing explicitly the integrals, that for each $0 \leq r < 1$ and for each $0 \leq \theta \leq 2\pi$:

a)

$$\frac{1}{2\pi} \int_0^{2\pi} \frac{1-r^2}{1-2r \cos(\theta-\phi) + r^2} d\phi = 1$$

Solution:

Let $D = \{(r, \theta) \mid 0 \leq r \leq 1, 0 \leq \theta \leq 2\pi\}$ be the unit disk and u the constant function equal to 1

$$\begin{aligned} u : D &\rightarrow \mathbb{R} \\ (r, \theta) &\mapsto u(r, \theta) \equiv 1. \end{aligned}$$

Then clearly u is the solution of the Dirichlet problem

$$\begin{cases} \nabla^2 u = 0, & \text{in } D \\ u(1, \theta) = 1, & 0 \leq \theta \leq 2\pi \end{cases}$$

and therefore for each $(r, \theta) \in [0, 1) \times [0, 2\pi]$ it must be represented by the Poisson integral formula:

$$\begin{aligned} u(r, \theta) &= \frac{1}{2\pi} \int_0^{2\pi} K(r, \theta, 1, \phi) u(1, \phi) d\phi \quad \rightsquigarrow \\ &\rightsquigarrow 1 = \frac{1}{2\pi} \int_0^{2\pi} \frac{1-r^2}{1-2r \cos(\theta-\phi) + r^2} d\phi. \end{aligned}$$

b)

$$\frac{1}{2\pi} \int_0^{2\pi} \frac{(1-r^2)(\cos^3(\phi)\sin(\phi) - \sin^3(\phi)\cos(\phi))}{1-2r\cos(\theta-\phi)+r^2} d\phi = \frac{r^3}{4} \sin(4\theta).$$

Solution:

We would like to do as we did before. That is, consider the function

$$f(\theta) = \cos^3(\theta)\sin(\theta) - \sin^3(\theta)\cos(\theta), \quad 0 \leq \theta \leq 2\pi.$$

We want to solve the Dirichlet problem on the unit disk with this boundary condition

$$\begin{cases} \nabla^2 u = 0, & \text{in } D \\ u(1, \theta) = f(\theta), & 0 \leq \theta \leq 2\pi \end{cases}$$

We can proceed by matching the coefficients in the Fourier series as did in the Exercises 1. and 2..

Even better, we can also recognise that the boundary condition

$$\cos^3(\theta)\sin(\theta) - \sin^3(\theta)\cos(\theta) = x^3y - xy^3$$

is already a well defined harmonic function on the whole plane

$$\nabla^2(x^3y - xy^3) = 6xy - 6xy \equiv 0.$$

So it must be itself the solution of the Dirichlet problem above.

Now we just need to manipulate it in polar coordinates to obtain

$$\begin{aligned} x^3y - xy^3 &= xy(x^2 - y^2) = r^3 \cos(\theta)\sin(\theta)(\cos^2(\theta) - \sin^2(\theta)) = \\ &= \frac{r^3}{2} \sin(2\theta)\cos(2\theta) = \frac{r^3}{4} \sin(4\theta), \end{aligned}$$

and finally using the Poisson integral formula we have proved that

$$\frac{1}{2\pi} \int_0^{2\pi} \frac{(1-r^2)(\cos^3(\phi)\sin(\phi) - \sin^3(\phi)\cos(\phi))}{1-2r\cos(\theta-\phi)+r^2} d\phi = \frac{r^3}{4} \sin(4\theta).$$