

Mathematical Foundations for Finance

Exercise sheet 11

Please hand in your solutions until Tuesday, 04/12/2018, 18:00 into your assistant's box next to HG G 53.2.

Exercise 11.1 Let (Ω, \mathcal{F}, P) be a probability space with a filtration $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$ satisfying the usual conditions. Assume that \mathcal{F}_0 is P -trivial and consider a Brownian motion W on this space.

- Prove that any continuous, adapted process H is predictable and locally bounded.
Hint 1: Recall that a process X is locally bounded if there exists a sequence of stopping times $(\tau_n)_{n \in \mathbb{N}}$ increasing to infinity such that each X^{τ_n} is uniformly bounded P -a.s.
- Prove that any predictable, locally bounded process H is an element of $L_{loc}^2(W)$.
Hint: We saw that $L_{loc}^2(M)$ can be characterized in a nice way when M is a continuous local martingale null at 0.
- Deduce that for any function $f : \mathbb{R} \rightarrow \mathbb{R}$ in C^1 , the stochastic integral $\int_0^\cdot f'(W_s) dW_s$ is a continuous local martingale.
- Conclude using Itô's formula that $f(W)$ for a given $f \in C^2$ is a continuous local martingale if and only if $\int_0^\cdot f''(W_s) ds = 0$.
*Hint 1: If M and N are local (P, \mathbb{F}) -martingales, then $M + N$ is a local (P, \mathbb{F}) -martingale.
Hint 2: For every continuous local martingale M null at 0 and with finite variation, we have that $M = 0$ P -a.s.*

Exercise 11.2 Let $W = (W_t)_{t \geq 0}$ be a Brownian motion with respect to a probability measure P and a filtration $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$. Using Itô's formula, decide for each of the following processes whether they are local (P, \mathbb{F}) -martingales or not. Which of them are even (P, \mathbb{F}) -martingales?

- $X_t^{(1)} := \exp\left(\frac{1}{2}\alpha^2 t\right) \cos(\alpha(W_t - \beta))$, $t \geq 0$, where $\alpha, \beta \in \mathbb{R}$.
Hint: For the martingale property of $X^{(1)}$, look first at $[0, T]$ for some $T > 0$.
- $X_t^{(2)} := \sin W_t - \cos W_t$, $t \geq 0$.
- $X_t^{(3)} := W_t^p - ptW_t$, $t \geq 0$, for $p \in \mathbb{N}$ with $p \geq 2$.
Hint: For any $T > 0$, $\sup_{0 \leq t \leq T} W_t$ has the same distribution as $|W_T|$ and so has $-\inf_{0 \leq s \leq T} W_s$.

Exercise 11.3 Let $W = (W_t)_{t \geq 0}$ be a Brownian motion with respect to some probability measure P and a filtration $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$. Use Itô's formula to write the following processes as stochastic integrals.

- $X_t^{(1)} = W_t^2$.
- $X_t^{(2)} = t^2 W_t^3$.
- $X_t^{(3)} = \exp(mt + \sigma W_t)$.
- $X_t^{(4)} = \cos(t + W_t)$.
- $X_t^{(5)} = \log(2 + \cos(W_t - t))$.
- Let X and Y be two continuous real-valued (P, \mathbb{F}) -semimartingales. Define the process $Z = XY$. Apply Itô's formula to Z and write it as a sum of stochastic integrals.