

**Multiple Choice.****Version D**

1. Let  $G$  be a group and  $H$  a subgroup of  $G$ . Which of the following statements are *always* true in this case?
  - (a) **(Right)** For  $G$  finite, the index of  $H$  in  $G$  divides the order of  $G$ .
  - (b) **(Wrong)**  $G$  contains a normal subgroup whose index is a prime number.
  - (c) **(Wrong)** If  $H$  is abelian and normal in  $G$  and  $G/H$  is abelian, then  $G$  is abelian.
  - (d) **(Right)** If the index of  $H$  in  $G$  is two, the subgroup  $H$  is a normal subgroup of  $G$ .
  - (e) **(Wrong)** For any two simple groups  $G_1, G_2$ , their direct sum  $G_1 \oplus G_2$  is also simple.

Recall: A group  $G$  is called *simple* if the only normal subgroups of  $G$  are  $\{1\}$  and  $G$ .

2. Let  $G$  be a group acting on a set  $X$  and let  $H$  be a subgroup of  $G$ . Which of the following statements are *always* true in this case?
  - (a) **(Right)** For any field  $k$ , the action of the group  $\mathrm{SL}_n(k)$  on  $k^n$  where a matrix  $A \in \mathrm{SL}_n(k)$  acts on a vector  $v \in k^n$  by multiplication from the left is a faithful action.
  - (b) **(Right)** If there exists  $x \in X$  with  $\mathrm{Stab}_G(x) = \{1\}$  then the action of  $G$  on  $X$  is faithful.
  - (c) **(Right)** If the action of  $H$  on  $X$  is transitive, so is the action of  $G$  on  $X$ .
  - (d) **(Wrong)** There is a map from the set of  $G$ -orbits to the set of  $H$ -orbits which sends  $Gx$  to  $Hx$ .
  - (e) **(Wrong)** The only action of the group  $G = \mathbb{Z}/2\mathbb{Z}$  on the set  $X = \{1, 2, 3\}$  is the trivial action  $gx = x$  for  $g \in G, x \in X$ .

Recall: An action of  $G$  on  $X$  is called *faithful* if the only element  $g \in G$  satisfying  $gx = x$  for all  $x \in X$  is the neutral element  $g = 1$ .

3.
  - (a) **(Right)** The group  $S_n$  has a subgroup of order  $m$  for each  $m = 1, 2, \dots, n$ .
  - (b) **(Right)** The 6-cycle  $\sigma = (123456) \in S_6$  has a decomposition into a product  $\sigma = \tau_1 \tau_2 \cdots \tau_m$  of transpositions  $\tau_1, \tau_2, \dots, \tau_m \in S_6$  and for each such decomposition the number  $m$  of transpositions is odd.
  - (c) **(Right)** The 6-cycle  $\sigma = (123456) \in S_6$  cannot be written as the product of 3 transpositions.
  - (d) **(Wrong)** The group  $A_5$  is the only simple subgroup of  $S_5$ .
  - (e) **(Wrong)** The subgroup  $G = \langle (12), (345) \rangle \subset S_5$  is isomorphic to  $S_6$ .

4. Let  $A$  be a commutative ring and  $I$  an ideal of  $A$ . Which of the following statements are then *always* true?

- (a) **(Wrong)** For all  $f, g \in A[X]$  of degree 3, we have that  $f + g$  also has degree 3.
- (b) **(Right)** If  $I$  is a prime ideal, then for  $a, b \in A/I$  with  $a \neq 0, b \neq 0$  we have  $ab \neq 0$ .
- (c) **(Wrong)** For any PID  $A$ , also  $A[X]$  is a PID.
- (d) **(Wrong)** The ideal  $I = (X)$  in  $A[X]$  is maximal.
- (e) **(Right)** If  $I \subsetneq A$  is a proper ideal, then there exists a prime ideal of  $A$  containing  $I$ .

Recall: We say that a polynomial  $f \in A[X]$  has degree  $m$  if  $f$  has the form  $f(X) = a_m X^m + a_{m-1} X^{m-1} + \dots + a_1 X + a_0$  with  $a_m \neq 0$ .

5. Let  $A$  and  $B$  be commutative rings and  $f : A \rightarrow B$  a ring homomorphism. Let  $I$  be an ideal of  $A$  and denote by  $p : A \rightarrow A/I$  the usual projection. Which of the following statements are then *always* true?

- (a) **(Right)** There exists a ring homomorphism  $g : A/I \rightarrow B$  satisfying  $f = g \circ p$  if and only if  $f$  vanishes on  $I$ .
- (b) **(Right)** If  $f \neq 0$  and  $B$  is an integral domain, then  $A/\ker(f)$  is an integral domain.
- (c) **(Wrong)** If  $f$  is surjective, then  $B$  is isomorphic to a subring of  $A$ .
- (d) **(Right)** For every  $b_1, b_2 \in B$  there exists a unique ring homomorphism  $h : A[X, Y] \rightarrow B$  sending  $X \mapsto b_1, Y \mapsto b_2$  and such that  $h|_A = f$ , where we use the natural inclusion  $A \subset A[X, Y]$ .
- (e) **(Right)** If  $J \subset B$  is a maximal ideal and  $f : A \rightarrow B$  is surjective, then  $f^{-1}(J)$  is a maximal ideal in  $A$ .

6. Let  $R = \mathbb{Z}/30\mathbb{Z}$ .

- (a) **(Right)** The natural map  $R \rightarrow (\mathbb{Z}/2\mathbb{Z}) \oplus (\mathbb{Z}/3\mathbb{Z}) \oplus (\mathbb{Z}/5\mathbb{Z})$  sending  $\bar{a}$  to  $(\bar{a}, \bar{a}, \bar{a})$  is an isomorphism.
- (b) **(Wrong)** There exist exactly 3 elements of order 3 in the group  $(R, +, 0)$ .
- (c) **(Right)** The ideal  $(\bar{5}) \subset R$  is maximal.
- (d) **(Wrong)** There exist exactly 3 distinct ring homomorphisms  $R \rightarrow \mathbb{Z}/3\mathbb{Z}$ .
- (e) **(Right)** Every ideal in  $R$  is a principal ideal.

7. (a) **(Wrong)** Every torsion-free  $\mathbb{Z}$ -module is free.

- (b) **(Right)** Every free  $\mathbb{Z}$ -module  $A$  with  $A \neq 0$  contains  $\mathbb{Z}$  as a submodule.
- (c) **(Wrong)** There are, up to isomorphism, 4 different abelian groups of 8 elements.
- (d) **(Right)** If  $A$  is a finitely generated  $\mathbb{Z}$ -module in which each element has finite order, then  $A$  is finite.
- (e) **(Right)** If  $A$  is a finitely generated  $\mathbb{Z}$ -module in which each nonzero element has infinite order, then  $A$  is free.

8. Let  $L/K$  be a field extension.

- (a) **(Right)** If  $\dim_K L = 17$  then the field extension is algebraic.
- (b) **(Wrong)** If  $f \in K[T]$  has two distinct roots in  $L$ , then  $f$  cannot be irreducible in  $K[T]$ .
- (c) **(Right)** If  $\alpha \in L \setminus \{0\}$  is transcendental over  $K$ , then also  $1/\alpha$  is transcendental.
- (d) **(Right)** If  $\alpha \in L$  is algebraic and  $L/K$  is of finite degree, then  $\deg(\text{irr}(\alpha; k))$  divides  $[L : K]$ .
- (e) **(Wrong)** For  $\alpha \in L$ , the field  $K(\alpha) \subset L$  is equal to the  $K$ -vector subspace of  $L$  spanned by  $1, \alpha, \alpha^2, \dots$ .

Recall: If  $\alpha \in L$  is algebraic over  $K$ , its minimal polynomial  $\text{irr}(\alpha; k)$  is the unique monic polynomial in  $K[X]$  of minimal degree among all nonzero polynomials in  $K[X]$  that vanish on  $\alpha$ .

9. (a) **(Right)** For every finite field  $\mathbb{F}$ , the characteristic  $\text{char}(\mathbb{F})$  divides the number of elements  $|\mathbb{F}|$  of  $\mathbb{F}$ .
- (b) **(Wrong)** If  $K$  is an extension of degree 2 of the finite field  $\mathbb{F}_4$ , then  $K$  is isomorphic to  $\mathbb{F}_8$ .
- (c) **(Wrong)** For each element  $a \in \mathbb{F}_q \setminus \{0\}$  one has  $a^q = 1$ .
- (d) **(Wrong)** The polynomial  $X^{121} - 1$  has exactly 10 distinct roots in  $\mathbb{F}_{11}$ .
- (e) **(Wrong)** If  $E$  is a finite field and  $F/E$  is an algebraic extension, then  $F$  is a finite field.

Recall: For a field  $K$  the characteristic  $\text{char}(K)$  is defined to be  $\text{char}(K) = 0$  if there exists no integer  $n \in \mathbb{Z}_{>0}$  with  $n \cdot 1_K = 0_K$ . Otherwise,  $\text{char}(K)$  is defined to be the minimal positive integer  $n \in \mathbb{Z}_{>0}$  with  $n \cdot 1_K = 0_K$ .

10. (a) **(Right)** The splitting field in  $\mathbb{C}$  of the  $\mathbb{Q}$ -polynomial  $X^2 - 2$  is  $\mathbb{Q}(\sqrt{2})$ .
- (b) **(Wrong)** The splitting field in  $\mathbb{C}$  of the  $\mathbb{Q}$ -polynomial  $X^2 + 5$  is  $\mathbb{Q}(\sqrt{5}, i)$ .
- (c) **(Wrong)** The splitting field in  $\mathbb{C}$  of the  $\mathbb{Q}$ -polynomial  $X^{17} - 1$  is isomorphic to  $\mathbb{Q}[X]/(X^{17} - 1)$ .
- (d) **(Wrong)** For any  $\alpha \in \mathbb{C}$  there exists a  $\mathbb{Q}$ -polynomial  $f(X) \in \mathbb{Q}[X]$  such that  $\mathbb{Q}(\alpha)$  is the splitting field in  $\mathbb{C}$  of  $f(X)$ .
- (e) **(Right)** The field  $\mathbb{Q}(\sqrt{7}, i)$  has degree 4 over  $\mathbb{Q}$ .
-