

# Mathematical Finance

## Exercise sheet 10

**Exercise 10.1** Consider a financial market  $S$ . Let

$$\mathcal{X}_1 = \{1 + \vartheta \bullet S : \vartheta \in \Theta_{\text{adm}}^1\}.$$

Show that  $\mathcal{X}_1$  satisfies the switching property: for any  $X \in \mathcal{X}_1$  and any strictly positive  $X' \in \mathcal{X}_1$ , a stopping time  $\tau$  and event  $A \in \mathcal{F}_\tau$ , we have that

$$\tilde{X} = \mathbb{1}_{\Omega \setminus A} X + \mathbb{1}_A \frac{X'_\tau \vee X_\tau}{X'_\tau} X_{\tau \wedge \cdot}.$$

belongs to  $\mathcal{X}_1$ .

**Solution 10.1** Let  $X, X' \in \mathcal{X}_1$ , with  $X' > 0$ . Then, we can find  $\vartheta, \vartheta' \in \Theta_{\text{adm}}^1$  such that  $X = 1 + \vartheta \bullet S$  and  $X' = 1 + \vartheta' \bullet S$ .

Now, note that

$$\hat{X} := 1 + \left[ \mathbb{1}_{[0, \tau]} \vartheta + (\mathbb{1}_{(\tau, T]} \mathbb{1}_{\Omega \setminus A} \vartheta) + \left( \mathbb{1}_{(\tau, T]} \mathbb{1}_A \frac{X_\tau}{X'_\tau} \vartheta' \right) \right] \bullet S = \tilde{X}.$$

Indeed, we can check that  $\hat{X} = X = \tilde{X}$  on  $\Omega \setminus A$  and on  $A$  if  $0 \leq t \leq \tau$ , while on  $A$  with  $t \geq \tau$ ,

$$\begin{aligned} \hat{X}_t &= 1 + \int_0^\tau \vartheta_s dS_s + \int_\tau^t \frac{X_\tau}{X'_\tau} \vartheta'_s dS_s \\ &= X_\tau \left( 1 + \frac{1}{X'_\tau} \int_\tau^t \vartheta'_s dS_s \right) \\ &= X_\tau \left( 1 + \frac{1}{X'_\tau} (X'_t - X'_\tau) \right) \\ &= \frac{X_\tau}{X'_\tau} X'_t = \tilde{X}_t. \end{aligned}$$

Moreover, note that the strategy  $\tilde{\vartheta} := \mathbb{1}_{[0, \tau]} \vartheta + (\mathbb{1}_{(\tau, T]} \mathbb{1}_{\Omega \setminus A} \vartheta) + \left( \mathbb{1}_{(\tau, T]} \mathbb{1}_A \frac{X_\tau}{X'_\tau} \vartheta' \right)$  is predictable, since each of  $\vartheta, \vartheta', \mathbb{1}_{[0, \tau]}, \mathbb{1}_{(\tau, T]} \mathbb{1}_{\Omega \setminus A}, \mathbb{1}_{(\tau, T]} \mathbb{1}_A, \mathbb{1}_{(\tau, T]} X_\tau, \mathbb{1}_{(\tau, T]} \frac{1}{X'_\tau}$  is predictable. The fact that  $\tilde{\vartheta} \in L(S)$  can be deduced easily from Theorem 1.29 in the lecture notes.

For the following questions, we consider a financial market satisfying (NFLVR), with a numéraire (which we set to be equal to 1) and a  $d$ -dimensional risky asset with discounted prices  $S_t$  taking values in  $D \subseteq \mathbb{R}^d$ .

We work under an ESM,  $Q$ , and we will work with polynomial models. Consider the following two models:

- The Black-Scholes model:

$$dS_t = S_t \sigma dW_t \quad (S_0 \in \mathbb{R}_{>0}^d),$$

where  $\sigma \in \mathbb{R}^{d \times d}$  is invertible, and  $W$  is a  $d$ -dimensional Brownian motion.  $S_t$  in the right-hand side is viewed as a diagonal matrix with entries  $S^i$ .

- The SABR model (in Bachelier form):

$$dS_t^1 = S_t^2 dW_t, \quad dS_t^2 = \alpha S_t^2 dB_t \quad (S_0^1, S_0^2 > 0),$$

for some parameter  $\alpha > 0$  and Brownian motions  $W, B$  with fixed correlation  $\rho \in [-1, 1]$ .  $S^2$  is the stochastic volatility, which we assume that we can trade through forward volatility contracts.

For these two models, solve the following exercises:

**Exercise 10.2** Show that  $S$  is a polynomial process, in other words, that for any  $s \leq t$  and any polynomial  $p$  of degree  $n$  we have

$$E[p(S_t) \mid \mathcal{F}_s] = q(S_s),$$

where  $q$  is a polynomial of degree  $\leq n$ , whose coefficients are functions of  $t - s$ .

**Solution 10.2** In the Black-Scholes, consider a monomial  $x_1^{k_1} x_2^{k_2} \dots x_d^{k_d}$ . We compute

$$\begin{aligned} \prod_{j=1}^d (S_t^j)^{k_j} &= \prod_{j=1}^d (S_0^j)^{k_j} \exp \left( \left( \sum_{j=1}^d k_j \sigma_j \right) B_t - \frac{t}{2} \sum_{j=1}^d k_j |\sigma_j|^2 \right) \\ &= \prod_{j=1}^d (S_s^j)^{k_j} \exp \left( \left( \sum_{j=1}^d k_j \sigma_j \right) (B_t - B_s) - \frac{t-s}{2} \sum_{j=1}^d k_j |\sigma_j|^2 \right), \end{aligned}$$

where  $\sigma_j$  are the rows of  $\sigma$ , i.e.  $\sigma_j = (\sigma_{j1}, \dots, \sigma_{jd})$ . This can be further decomposed as

$$\prod_{j=1}^d (S_t^j)^{k_j} = \prod_{j=1}^d (S_s^j)^{k_j} \exp \left( \left( \sum_{j=1}^d k_j \sigma_j \right) (B_t - B_s) - \frac{t-s}{2} \left| \sum_{j=1}^d k_j \sigma_j \right|^2 \right) \exp \left( \frac{t-s}{2} \left( \left| \sum_{j=1}^d k_j \sigma_j \right|^2 - \sum_{j=1}^d k_j |\sigma_j|^2 \right) \right).$$

Since the middle term is a martingale (by Novikov), we obtain

$$E \left[ \prod_{j=1}^d (S_t^j)^{k_j} \mid \mathcal{F}_s \right] = \prod_{j=1}^d (S_s^j)^{k_j} \exp \left( \kappa \frac{t-s}{2} \right),$$

with  $\kappa = \left( \left| \sum_{j=1}^d k_j \sigma_j \right|^2 - \sum_{j=1}^d k_j |\sigma_j|^2 \right)$ . We observe that  $S$  is indeed a polynomial process.

For the SABR model, we consider the monomial  $x_1^{k_1} x_2^{k_2}$ , and fix some time  $t \geq 0$ . We make the ansatz that we can find a martingale of the form

$$M_s = \sum_{j=0}^{k_1} c_j(t-s) (S_s^1)^j (S_s^2)^{k_1+k_2-j},$$

where  $c_j$  are some deterministic, differentiable functions of time such that  $c_{k_1}(0) = 1$  and  $c_j(0) = 0$  for other values of  $j$ . By Itô's formula, we obtain

$$\begin{aligned}
dM_s &= \sum_{j=1}^{k_1} j c_j(t-s) (S_s^1)^{j-1} (S_s^2)^{k_1+k_2-j+1} dW_t + \sum_{j=0}^{k_1} \alpha(k_1+k_2-j) c_j(t-s) (S_s^1)^j (S_s^2)^{k_1+k_2-j} dB_t \\
&\quad - \sum_{j=0}^{k_1} c'_j(t-s) (S_s^1)^j (S_s^2)^{k_1+k_2-j} ds + \frac{1}{2} \sum_{j=2}^{k_1} j(j-1) c_j(t-s) (S_s^1)^{j-2} (S_s^2)^{k_1+k_2-j+2} ds \\
&\quad + \frac{1}{2} \sum_{j=0}^{k_1} \alpha^2(k_1+k_2-j)(k_1+k_2-j-1) c_j(t-s) (S_s^1)^j (S_s^2)^{k_1+k_2-j} ds \\
&\quad + \rho \sum_{j=1}^{k_1} \alpha j(k_1+k_2-j) c_j(t-s) (S_s^1)^{j-1} (S_s^2)^{k_1+k_2-j+1} ds.
\end{aligned}$$

Collecting the coefficients, we obtain the following system of ODEs:

$$c'_j(u) = \frac{(j+2)(j+1)}{2} c_{j+2}(u) + \frac{\alpha^2(k_1+k_2-j)(k_1+k_2-j-1)}{2} c_j(u) + \rho \alpha(j+1)(k_1+k_2-j-1) c_{j+1}(u),$$

where we set  $c_{k_1+1} = c_{k_1+2} = 0$ . Letting  $\mathbf{c} = (c_0, \dots, c_{k_1})^T$ , we can write this equation in the form

$$\mathbf{c}' = B\mathbf{c}$$

for a certain matrix  $B$  with the coefficients of the equation, and therefore we obtain

$$\mathbf{c}(t) = \exp(tB)\mathbf{c}_0$$

where  $\mathbf{c}_0 = (0, 0, \dots, 0, 1)^T$ . In conclusion, we obtain that  $M_s$  is a local martingale, where the coefficients  $\mathbf{c}$  are given above. One can show that the moments of  $S$  are finite, which implies that  $M$  is even a true martingale. Therefore, we obtain that

$$E[(S_t^1)^{k_1} (S_t^2)^{k_2} | \mathcal{F}_s] = E[M_t | \mathcal{F}_s] = M_s = \sum_{j=0}^{k_1} c_j(t-s) (S_s^1)^j (S_s^2)^{k_1+k_2-j}.$$

Thus,  $S$  is a polynomial model.

**Exercise 10.3** Find the delta hedge for a payoff of the form  $p(S_t)$ , for  $p$  a polynomial.

**Solution 10.3** For the Black-Scholes model, we use Itô's formula to differentiate

$$\begin{aligned}
d \left( \prod_{j=1}^d (S_s^j)^{k_j} \exp \left( \kappa \frac{t-s}{2} \right) \right) &= \exp \left( \kappa \frac{t-s}{2} \right) \prod_{j=1}^d (S_s^j)^{k_j} \sum_{j=1}^d k_j \sigma_j dW_s \\
&= \exp \left( \kappa \frac{t-s}{2} \right) \sum_{j=1}^d \left( k_j (S_s^j)^{k_j-1} \prod_{i \neq j} (S_s^i)^{k_i} \right) dS_s^j.
\end{aligned}$$

Thus, the delta hedge is given by the strategy  $\vartheta = (\vartheta^1, \dots, \vartheta^d)^T$ , with

$$\vartheta_s^j = k_j \exp \left( \kappa \frac{t-s}{2} \right) (S_s^j)^{k_j-1} \prod_{i \neq j} (S_s^i)^{k_i},$$

where  $\vartheta^j = 0$  if  $k_j = 0$ . The initial value is

$$\varphi_0 = \prod_{j=1}^d (S_0^j)^{k_j} \exp\left(\frac{\kappa t}{2}\right).$$

Likewise, for the SABR model, we obtain

$$dM_s = \left( \sum_{j=0}^{k_1} j c_j(t-s) (S_s^1)^{j-1} (S_s^2)^{k_1+k_2-j} \right) dS_s^1 + \left( \sum_{j=0}^{k_1} (k_1+k_2-j) c_j(t-s) (S_s^1)^j (S_s^2)^{k_1+k_2-j-1} \right) dS_s^2,$$

so that the delta hedge is given by

$$\begin{aligned} \vartheta_s^1 &= \sum_{j=0}^{k_1} j c_j(t-s) (S_s^1)^{j-1} (S_s^2)^{k_1+k_2-j}, \\ \vartheta_s^2 &= \sum_{j=0}^{k_1} (k_1+k_2-j) c_j(t-s) (S_s^1)^j (S_s^2)^{k_1+k_2-j-1}, \end{aligned}$$

as well as  $\varphi_0 = M_0$ .

**Exercise 10.4** Note that the models under consideration are Markovian. Define and compute the transition semigroup  $(P_t)_{t \geq 0}$  (under  $Q$ ), as it acts on the set of real polynomials  $\text{Pol}(\mathbb{R}^d)$  ( $\mathbb{R}^2$  in the case of the SABR model).

Show that in this setting,

$$P_{t-s} f(X_s) = (\nabla P_{t-s} f(S) \bullet S)_s + P_t(x), \quad Q^x\text{-a.s.},$$

where  $Q^x$  is the law of  $S$  started from a given point  $x$ .

Show that this equality also holds  $P$ -almost surely, where  $P$  is the historical measure.

**Solution 10.4** The transition semigroup  $P_t$  acts on  $\text{Pol}(\mathbb{R}^d)$  by

$$P_t(f) = g,$$

where  $f$  is some polynomial and  $g$  is the unique polynomial such that

$$E[f(S_t) \mid \mathcal{F}_0] = g(S_0),$$

or in other words,

$$E_x[f(S_t)] = g(x).$$

Thanks to the calculations above, we can compute  $P_t$  for monomials (and thus all polynomials, by linearity):

$$P_t \left( \prod_{j=1}^d x_j^{k_j} \right) = \exp\left(\frac{\kappa t}{2}\right) \prod_{j=1}^d x_j^{k_j},$$

for the Black-Scholes model (note that  $\kappa$  depends on the coefficients  $k_j$ ).

On the other hand, for the SABR model, we have

$$P_t(x_1^{k_1} x_2^{k_2}) = \sum_{j=0}^{k_1} c_j(t) (x_1)^j (x_2)^{k_1+k_2-j}.$$

From our previous considerations, in order to show the equality it is enough to check that

$$\nabla P_{t-} f(S) = \vartheta.$$

This is a simple calculation: in the Black-Scholes model, and for  $f$  a monomial as above,

$$\begin{aligned} \partial_j P_{t-s} f(S_s) &= \partial_j \left( \exp\left(\frac{\kappa(t-s)}{2}\right) \prod_{j=1}^d x_j^{k_j} \right) \Big|_{x=S_s} \\ &= k_j \exp\left(\frac{\kappa(t-s)}{2}\right) (S_s^j)^{k_j-1} \prod_{i \neq j} x_i^{k_i} \\ &= k_j \exp\left(\frac{\kappa(t-s)}{2}\right) (S_s^j)^{k_j-1} \prod_{i \neq j} (S_s^i)^{k_i} \\ &= \vartheta_s^j. \end{aligned}$$

Finally, the equality also holds  $P$ -a.s., since stochastic integrals are preserved under equivalent measures, and likewise, almost sure equality is also preserved.

### Exercise 10.5 (Python)

Implement the discretised delta hedge from exercise 3, for the payoff  $H = (S_T)^3$ . Compute the error between the hedge and the payoff, as well as the difference between the payoff and a hedging strategy that only trades in  $S$ , but not  $Y$ .