

Exercise Sheet 1

1. a.) The Möbius function $\mu : \mathbb{N} \rightarrow \{\pm 1, 0\}$ is defined as

$$\mu(n) := \begin{cases} 0, & \text{if } n \text{ is not squarefree,} \\ (-1)^l, & \text{if } n = p_1 \cdots p_l \text{ is a product of distinct primes.} \end{cases}$$

Show that

$$\sum_{d|n} \mu(d) = \begin{cases} 1, & \text{if } n = 1, \\ 0, & \text{otherwise.} \end{cases}$$

Hint: Use the inclusion-exclusion principle.

b.) Recall that the Euler totient function $\varphi : \mathbb{N} \rightarrow \mathbb{N}$ is defined as

$$\varphi(n) := |\{1 \leq k \leq n : (k, n) = 1\}|.$$

Show that for all $n \geq 1$ we have

$$\varphi(n) = n \sum_{d|n} \frac{\mu(d)}{d}.$$

2. Given two functions $f, g : \mathbb{N} \rightarrow \mathbb{C}$ we can define their Dirichlet convolution $f * g : \mathbb{N} \rightarrow \mathbb{C}$ via

$$(f * g)(n) := \sum_{d|n} f(d) g\left(\frac{n}{d}\right).$$

a.) Show that for all $f, g, h : \mathbb{N} \rightarrow \mathbb{C}$ we have

$$f * g = g * f \quad \text{and} \quad (f * g) * h = f * (g * h).$$

b.) (Möbius inversion formula). Let $f : \mathbb{N} \rightarrow \mathbb{C}$ be any function. Show that

$$g(n) = \sum_{d|n} f(d) \text{ for all } n \in \mathbb{N} \quad \Leftrightarrow \quad f(n) = \sum_{d|n} g(d) \mu\left(\frac{n}{d}\right) \text{ for all } n \in \mathbb{N}.$$

c.) Show that for all $n \in \mathbb{N}$ we have

$$n = \sum_{d|n} \varphi(d).$$

Bitte wenden!

d.) Let $f, g : \mathbb{N} \rightarrow \mathbb{C}$ be functions, and let $s \in \mathbb{C}$ be such that the associated Dirichlet series

$$F(s) := \sum_{n \geq 1} \frac{f(n)}{n^s}, \quad G(s) := \sum_{n \geq 1} \frac{g(n)}{n^s}$$

converge absolutely. Show that the Dirichlet series associated to $f * g$ satisfies

$$\sum_{n \geq 1} \frac{(f * g)(n)}{n^s} = F(s)G(s).$$

3. For a given integer $N \geq 1$, consider the probability space

$$\Omega_N := \{1, \dots, N\}$$

and let X_N, Y_N be independent random variables which are uniformly distributed on Ω_N . The goal of this exercise is to prove that

$$\mathbb{P}[(X_N, Y_N) = 1] \rightarrow \frac{6}{\pi^2}$$

as $N \rightarrow \infty$.

a.) Prove that for any $s \in \mathbb{C}$ with $\Re s > 1$ we have

$$\sum_{n \geq 1} \frac{\mu(n)}{n^s} = \frac{1}{\zeta(s)}.$$

b.) Prove that

$$\mathbb{P}[(X_N, Y_N) = 1] \rightarrow \frac{1}{\zeta(2)}.$$

c.) Show that, in order to prove that $\zeta(2) = \frac{\pi^2}{6}$, it suffices to show that

$$\sum_{n \geq 1} \frac{1}{(2n-1)^2} = \frac{\pi^2}{8}.$$

d.) Prove that

$$I := \int_0^1 \int_0^1 \frac{1}{1-x^2y^2} dx dy = \sum_{n \geq 1} \frac{1}{(2n-1)^2}.$$

e.) Compute I , e.g. by substituting

$$u := \arccos \left(\sqrt{\frac{1-x^2}{1-x^2y^2}} \right), \quad v := \arccos \left(\sqrt{\frac{1-y^2}{1-x^2y^2}} \right),$$

so that $x = \frac{\sin u}{\cos v}$ and $y = \frac{\sin v}{\cos u}$.