

Exercise Sheet 7

1. Jacobi fields in space forms

Let M be a space form with curvature $\kappa \in \mathbb{R}$. Furthermore, let $c: \mathbb{R} \rightarrow M$ be a geodesic which is parametrized by arc length and $N_0 \in TM_{c(0)}$ with $|N_0| = 1$, $\langle N_0, \dot{c}(0) \rangle = 0$. Determine the Jacobi field Y along c with starting conditions $Y(0) = aN_0$ and $\dot{Y}(0) = bN_0$ for $a, b \in \mathbb{R}$.

2. Trace of a symmetric bilinear form

Let $(V, \langle \cdot, \cdot \rangle)$ be a m -dimensional Euclidean space and let $r: V \times V \rightarrow \mathbb{R}$ be a symmetric bilinear form. Furthermore, let $S^{m-1} = \{v \in V : |v| = 1\}$ be the unit sphere. Prove that

$$\int_{S^{m-1}} r(v, v) \, d\text{vol}^{S^{m-1}} = \frac{\text{vol}(S^{m-1})}{m} \text{tr}(r) = \omega_m \text{tr}(r),$$

where $d\text{vol}^{S^{m-1}}$ denotes the induced volume on S^{m-1} and ω_m is the volume of the m -dimensional unit ball.

3. Small balls and scalar curvature

Let p be a point in the m -dimensional Riemannian manifold (M, g) . The goal is to prove the following Taylor expansion of the volume of the ball $B_r(p)$ as a function of r :

$$\text{vol}(B_r(p)) = \omega_m r^m \left(1 - \frac{1}{6(m+2)} \text{scal}(p)r^2 + \mathcal{O}(r^3) \right).$$

- a) Let $v \in TM_p$ with $|v| = 1$, define the geodesic $c(t) := \exp_p(tv)$ and let $e_1 = v, e_2, \dots, e_m \in TM_p$ be an orthonormal basis. Consider the Jacobi fields Y_i along c with $Y_i(0) = 0$ and $\dot{Y}_i(0) = e_i$ for $i = 2, \dots, m$. Show that the volume distortion factor of \exp_p at tv is given by

$$J(v, t) := \sqrt{\det(\langle T_{tv}e_i, T_{tv}e_j \rangle)} = t^{-(m-1)} \sqrt{\det(\langle Y_i, Y_j \rangle)},$$

where $T_{tv} := (d\exp_p)_{tv}$.

- b) Let E_2, \dots, E_m be parallel vector fields along c with $E_i(0) = e_i$. Then the Taylor expansion of Y_i is

$$Y_i(t) = tE_i - \sum_{k=2}^m \left(\frac{t^3}{6} R(e_i, v, e_k, v) + \mathcal{O}(t^4) \right) E_k.$$

- c) Conclude that $J(v, t) = 1 - \frac{t^2}{6} \text{ric}(v, v) + \mathcal{O}(t^4)$.
Hint: Use $\det(I_m + \epsilon A) = 1 + \epsilon \text{tr}(A) + \mathcal{O}(\epsilon^2)$.
- d) Prove the above formula for $\text{vol}(B_r(p))$.

Submission: until 12:15, April 14th, by email at the following address:
`tommaso.goldhirsch@math.ethz.ch`