

Problem set 3

Let H be a homology theory.

1. Prove from the axioms that $H_i(\emptyset) = 0$ for all i and that $H_i(X, X) = 0$ for all i and all spaces X .
2. Check that the long exact sequence for reduced homology continues to hold for pairs of the type (X, \emptyset) , including the case $X = \emptyset$.
3. Let $A \subset X$ be a non-empty subset and assume that $\tilde{H}_*(A) = 0$ (that is, A is acyclic). Prove that $H_*(X, A) \cong \tilde{H}_*(X)$.
4. Define the *unreduced suspension* ΣX of a space X to be the quotient space of $[0, 1] \times X$ obtained by identifying $\{0\} \times X$ and $\{1\} \times X$ to points. Show that there is a natural isomorphism

$$\tilde{H}_i(X) \xrightarrow{\cong} \tilde{H}_{i+1}(\Sigma X).$$

Here *natural* means that for a map $f: X \rightarrow Y$, and its suspension $\Sigma f: \Sigma X \rightarrow \Sigma Y$ the following diagram commutes:

$$\begin{array}{ccc} \tilde{H}_i(X) & \xrightarrow{\cong} & \tilde{H}_{i+1}(\Sigma X) \\ \downarrow f_* & & \downarrow (\Sigma f)_* \\ \tilde{H}_i(Y) & \xrightarrow{\cong} & \tilde{H}_{i+1}(\Sigma Y). \end{array}$$

Hint: Consider the two cones $C_+X := \{[t, x] \in \Sigma X \mid t \geq \frac{1}{2}\}$ and $C_-X := \{[t, x] \in \Sigma X \mid t \leq \frac{1}{2}\}$.

5. Let X and Y be topological spaces. Let $x_0 \in X$ and $y_0 \in Y$ be points. Assume that x has a closed neighborhood N in X , of which $\{x_0\}$ is a strong deformation retract. Recall the wedge $X \vee Y$ of X and Y we defined in Problem Set 2, Exercise 5. Show that the inclusion maps induce isomorphisms

$$\tilde{H}_i(X) \oplus \tilde{H}_i(Y) \xrightarrow{\cong} \tilde{H}_i(X \vee Y),$$

whose inverse is induced by the projections of $X \vee Y$ to X and Y .

6. Let H be a “theory” satisfying axioms 1-4 of a homology theory, but not necessarily axiom 5. Show that

$$(i_X)_* \oplus (i_Y)_* : H_p(X) \oplus H_p(Y) \rightarrow H_p(X \sqcup Y)$$

is an isomorphism for all spaces X, Y and for all $p \in \mathbb{Z}$, where $i_X : X \hookrightarrow X \sqcup Y$, $i_Y : Y \hookrightarrow X \sqcup Y$ denote the inclusions into the disjoint union.

Hints.

- (a) Consider the long exact sequence of the pair $(X \sqcup Y, X)$.
 (b) Consider the excision $(Y, \emptyset) = ((X \sqcup Y) \setminus X, X \setminus X) \xrightarrow{k} (X \sqcup Y, X)$ and the resulting isomorphism $k_* : H_*(Y) \xrightarrow{\cong} H_*(X \sqcup Y, X)$.
 (c) Note that the following diagram commutes:

$$\begin{array}{ccc} (Y, \emptyset) & \xrightarrow{k} & (X \sqcup Y, X) \\ & \searrow i_Y & \nearrow j \\ & X \sqcup Y & \end{array}$$

- (d) Deduce that in the long exact sequence

$$\cdots \rightarrow H_p(X) \xrightarrow{(i_X)_*} H_p(X \sqcup Y) \xrightarrow{j_*} H_p(X \sqcup Y, X) \rightarrow \cdots$$

all maps j_* are surjective, and that thus the sequence gives rise to short exact sequences

$$0 \rightarrow H_p(X) \xrightarrow{(i_X)_*} H_p(X \sqcup Y) \xrightarrow{j_*} H_p(X \sqcup Y, X) \rightarrow 0.$$

- (e) Find a right inverse for j_* to show that these short exact sequences split.