

EXERCISE SHEET 2

Exercise 1.(Identity Neighborhoods Generate Connected Groups):

Let G be a connected topological group, $U \subset G$ an open neighborhood of the identity and $U^n := \{g_1 \cdots g_n \mid g_1, \dots, g_n \in U\}$. Show that $G = \bigcup_{n=1}^{\infty} U^n$.

Hint: You may assume that $g^{-1} \in U$ for every $g \in U$. Why?

Exercise 2.(Transitive Group Actions):

Let G be a topological group, X a topological space and $\mu : G \times X \rightarrow X$ a continuous transitive group action, i.e. for any two $x, y \in X$ there is $g \in G$ such that $\mu(g, x) = g \cdot x = y$.

- Show that if G is compact then X is compact.
- Show that if G is connected then X is connected.

Exercise 3.(Examples of Haar Measures):

- Let us consider the *three-dimensional Heisenberg group* $H = \mathbb{R} \rtimes_{\eta} \mathbb{R}^2$, where $\eta : \mathbb{R} \rightarrow \text{Aut}(\mathbb{R}^2)$ is defined by

$$\eta(x) \begin{pmatrix} y \\ z \end{pmatrix} = \begin{pmatrix} y \\ z + xy \end{pmatrix},$$

for all $x, y, z \in \mathbb{R}$. Thus the group operation is given by

$$(x_1, y_1, z_1) * (x_2, y_2, z_2) = (x_1 + x_2, y_1 + y_2, z_1 + z_2 + x_1 y_2)$$

and it is easy to see that it can be identified with the matrix group

$$H \cong \left\{ \begin{pmatrix} 1 & x & z \\ 0 & 1 & y \\ 0 & 0 & 1 \end{pmatrix} : x, y, z \in \mathbb{R} \right\}$$

Verify that the Lebesgue measure is the Haar measure of $\mathbb{R} \rtimes_{\eta} \mathbb{R}^2$ and that the group is unimodular.

- Let

$$P = \left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} : a, b \in \mathbb{R}, a \neq 0 \right\}.$$

Show that $\frac{da}{a^2} db$ is the left Haar measure and $da db$ is the right Haar measure. In particular, P is *not* unimodular.

- c) Let $G := \mathrm{GL}_n(\mathbb{R}) \subseteq \mathbb{R}^{n^2}$ denote the group of invertible matrices over \mathbb{R} . Let λ_{n^2} denote the Lebesgue measure on \mathbb{R}^{n^2} . Prove that

$$dm(x) := |\det x|^{-n} d\lambda_{n^2}(x)$$

defines a bi-invariant (i.e. left- and right-invariant) Haar measure on G .

- d) Let $G = \mathrm{SL}_n(\mathbb{R})$ denote the group of matrices of determinant 1 in $\mathbb{R}^{n \times n}$. For a Borel subset $B \subseteq \mathrm{SL}_n(\mathbb{R})$ define

$$m(B) := \lambda_{n^2}(\{tg; g \in B, t \in [0, 1]\}).$$

Show that m is a well-defined bi-invariant Haar measure on $\mathrm{SL}_n(\mathbb{R})$.

- e) Let G denote the $ax + b$ group defined as

$$G = \left\{ \begin{pmatrix} a & b \\ & 1 \end{pmatrix}; a \in \mathbb{R}^\times, b \in \mathbb{R} \right\}$$

Note that every element in G can be written in a unique fashion as a product of the form:

$$\begin{pmatrix} a & b \\ & 1 \end{pmatrix} = \begin{pmatrix} \alpha & \\ & 1 \end{pmatrix} \begin{pmatrix} 1 & \beta \\ & 1 \end{pmatrix}$$

where $\alpha \in \mathbb{R}^\times$ and $\beta \in \mathbb{R}$, which yields a coordinate system $\mathbb{R}^\times \times \mathbb{R} \leftrightarrow G$. Prove that

$$dm(\alpha, \beta) = \frac{1}{|\alpha|} d\alpha d\beta$$

defines a left Haar measure on G . Calculate $\Delta_G(\alpha, \beta)$ for $\alpha \in \mathbb{R}^\times$ and $\beta \in \mathbb{R}$.

Exercise 4. (Haar Measure and Transitive Actions):

Let G be a locally compact Hausdorff group and let X be a topological space. Suppose that G acts on X continuously and transitively. Let $o \in X$, and denote $\pi: G \rightarrow X, g \mapsto g \cdot o$. Further, let

$$H := \mathrm{Stab}(o) = \{h \in G | h \cdot o = o\}$$

be the stabilizer of o .

Suppose there is a continuous section $\sigma: X \rightarrow G$ of π , i.e. $\pi \circ \sigma = \mathrm{Id}_X$.

- a) Show that $\psi: X \times H \rightarrow G, (x, h) \mapsto \sigma(x)h$ is a homeomorphism.

Hint: Find a continuous inverse!

- b) Suppose there is a (left) Haar measure ν on H and suppose there is a left G -invariant Borel regular measure λ on X .

Show that the push-forward measure $\psi_*(\lambda \otimes \nu)$ is a (left) Haar measure on G .

- c) Find a Haar measure on $\text{Iso}(\mathbb{R}^2)$.

Exercise 5. ($\text{Aut}(\mathbb{R}^n, +) \cong \text{GL}(n, \mathbb{R})$):

For a topological group G , we denote by $\text{Aut}(G)$ the group of bijective, continuous homomorphisms of G with continuous inverse. Consider the locally compact Hausdorff group $G = (\mathbb{R}^n, +)$ where $n \in \mathbb{N}_0$.

- a) Show that $\text{Aut}(G)$, i.e. the group of bijective homomorphisms which are homeomorphisms as well, is given by $\text{GL}_n(\mathbb{R})$.
- b) Show that $\text{mod} : \text{Aut}(G) \rightarrow \mathbb{R}_{>0}$ is given by $\alpha \mapsto |\det \alpha|^{-1}$.
- c) Prove that there exists a discontinuous, bijective homomorphism from the additive group $(\mathbb{R}, +)$ to itself.

Exercise 6. (Iterated Quotient Measures):

Let G be a locally compact Hausdorff group. Show that if $H_1 \leq H_2 \leq G$ are closed subgroups and H_1, H_2, G are all unimodular then there exist invariant measures dx, dy, dz on $G/H_1, G/H_2$ and H_2/H_1 respectively such that

$$\int_{G/H_1} f(x) dx = \int_{G/H_2} \left(\int_{H_2/H_1} f(yz) dz \right) dy$$

for all $f \in C_c(G/H_1)$.

Exercise 7. (No $\text{SL}_2(\mathbb{R})$ -invariant Measure on $\text{SL}_2(\mathbb{R})/P$):

Let $G = \text{SL}_2(\mathbb{R})$ and P be the subgroup of upper triangular matrices. Show directly that there is no (non-trivial) finite G -invariant measure on G/P .

Hint: Identify $G/P \cong \mathbb{S}^1 \cong \mathbb{R} \cup \{\infty\}$ with the unit circle and consider a rotation

$$\begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$$

and a translation

$$\begin{pmatrix} 1 & t \\ 0 & 1 \end{pmatrix}.$$

Due date: Thursday, 15/10/2020

Please, upload your solution via the SAM upload tool.

In order to access the website you will need a NETHZ-account and you will have to be connected to the ETH-network. From outside the ETH network you can connect to the ETH network via VPN. Here are instructions on how to do that.

Make sure that your solution is **one PDF file** and that its **file name** is formatted in the following way:

solution_<number exercise sheet>_<last name>_<first name>.pdf

For example: If your first name is Alice, your last name is Miller, and you want to hand-in your solution to Exercise Sheet 2, then you will have to upload your solution as **one** PDF file with the following file name:

solution_2_Miller_Alice.pdf

Solutions that fail to comply with the above requirements will be ignored.