

## EXERCISE SHEET 6

### Exercise 1. (Quotients of Lie groups):

Let  $G$  be a Lie group and let  $K \leq G$  be a closed normal subgroup.

Show that  $G/K$  can be equipped with a Lie group structure such that the quotient map  $\pi: G \rightarrow G/K$  is a surjective Lie group homomorphism with kernel  $K$ .

### Exercise 2. (Joint eigenvectors):

Let  $G$  be a connected Lie group and let  $\pi: G \rightarrow GL(V)$  be a finite-dimensional complex representation.

A *joint eigenvector* of  $\{\pi(g) : g \in G\}$  is a vector  $v \in V$  such that there is a smooth homomorphism  $\chi: G \rightarrow \mathbb{C}$  with  $\pi(g)v = \chi(g) \cdot v$  for all  $g \in G$ . Similarly, a *joint eigenvector* of  $\{d_e\pi(X) : X \in \mathfrak{g}\}$  is a vector  $v \in V$  such that there is a linear functional  $\lambda: \mathfrak{g} \rightarrow \mathbb{C}$  with  $d_e\pi(X)v = \lambda(X) \cdot v$  for all  $X \in \mathfrak{g}$ .

Show that a vector  $v \in V$  is a joint eigenvector of  $\{d_e\pi(X) : X \in \mathfrak{g}\}$  if and only if it is a joint eigenvector of  $\{\pi(g) : g \in G\}$ . Moreover, show that  $\chi(\exp(X)) = e^{\lambda(X)}$  for all  $X \in \mathfrak{g}$  (with  $\chi: G \rightarrow \mathbb{C}$  and  $\lambda: \mathfrak{g} \rightarrow \mathbb{C}$  as above).

### Exercise 3. (Isomorphism theorems for Lie algebras):

Let  $\mathfrak{g}$  be a Lie algebra.

- a) Let  $\mathfrak{h} \trianglelefteq \mathfrak{g}$  be an ideal. Show that

$$[X + \mathfrak{h}, Y + \mathfrak{h}] := [X, Y] + \mathfrak{h}$$

defines a Lie algebra structure on  $\mathfrak{g}/\mathfrak{h}$ .

- b) Show that if  $\varphi: \mathfrak{g} \rightarrow \mathfrak{h}$  is a Lie algebra homomorphism then

$$\mathfrak{g}/\ker\varphi \cong \text{im}\varphi$$

as Lie algebras.

- c) Let  $\mathfrak{h} \subseteq \mathfrak{I}$  be ideals of  $\mathfrak{g}$ . Show that

$$\mathfrak{I}/\mathfrak{h} \trianglelefteq \mathfrak{g}/\mathfrak{h} \quad \text{and} \quad (\mathfrak{g}/\mathfrak{h})/(\mathfrak{I}/\mathfrak{h}) \cong \mathfrak{g}/\mathfrak{I}.$$

- d) Let  $\mathfrak{h}$  and  $\mathfrak{I}$  be ideals of  $\mathfrak{g}$ . Show that  $\mathfrak{h} + \mathfrak{I}$  and  $\mathfrak{h} \cap \mathfrak{I}$  are ideals in  $\mathfrak{g}$ , and that

$$\mathfrak{h}/(\mathfrak{h} \cap \mathfrak{I}) \cong (\mathfrak{h} + \mathfrak{I})/\mathfrak{I}.$$

**Exercise 4. (Solvable Lie algebras):**

- Show that Lie subalgebras and homomorphic images of solvable Lie algebras are solvable.
- Show that if  $\mathfrak{h}$  and  $I$  are solvable ideals of a Lie algebra  $\mathfrak{g}$  then  $\mathfrak{h}+I$  is a solvable ideal.  
Hint: Use exercise 3 d)).
- Deduce that every Lie algebra contains a unique maximal solvable ideal.

**Exercise 5. (Weight spaces and ideals):**

Let  $\mathfrak{g}$  be a Lie algebra, let  $\mathfrak{h} \trianglelefteq \mathfrak{g}$  be an ideal and let  $\pi: \mathfrak{g} \rightarrow \mathfrak{gl}(V)$  a finite-dimensional complex representation. For a given linear functional  $\lambda: \mathfrak{h} \rightarrow \mathbb{C}$  consider its weight space

$$V_{\lambda}^{\mathfrak{h}} := \{v \in V \mid \pi(X)v = \lambda(X)v \quad \forall X \in \mathfrak{h}\}.$$

Show that every weight space  $V_{\lambda}^{\mathfrak{h}}$  is invariant under  $\pi(\mathfrak{g})$ , i.e.  $\pi(Y)V_{\lambda}^{\mathfrak{h}} \subseteq V_{\lambda}^{\mathfrak{h}}$  for every  $\lambda \in \mathfrak{h}^*$ ,  $Y \in \mathfrak{g}$ .

**Exercise 6. (Lie's theorem for Lie algebras):**

Let  $\mathfrak{g}$  be a solvable Lie algebra and let  $\rho: \mathfrak{g} \rightarrow \mathfrak{gl}(V)$  be a finite-dimensional complex representation.

Show that  $\rho(\mathfrak{g})$  stabilizes a flag  $V = V_0 \supseteq V_1 \supseteq \dots \supseteq V_n = 0$ , with  $\text{codim } V_i = i$ , i.e.  $\rho(X)V_i \subseteq V_i$  for every  $X \in \mathfrak{g}$ ,  $i = 1, \dots, n$ .

Hint: Use exercise 5.

**Due date:** Thursday, 10/12/2020

**Please, upload your solution via the SAM upload tool.**

In order to access the website you will need a NETHZ-account and you will have to be connected to the ETH-network. From outside the ETH network you can connect to the ETH network via VPN. Here are instructions on how to do that.

Make sure that your solution is **one PDF file** and that its **file name** is formatted in the following way:

solution\_<number exercise sheet>\_<last name>\_<first name>.pdf

**For example:** If your first name is Alice, your last name is Miller, and you want to hand-in your solution to Exercise Sheet 2, then you will have to upload your solution as **one** PDF file with the following file name:

solution\_2\_Miller\_Alice.pdf

**Solutions that fail to comply with the above requirements will be ignored.**