

# Mathematik III

## Final Exam

---

Identifying number: 001

---

- Put your student identity card on the desk. **Do not open this booklet** until you are so instructed.
- The exam is pseudonymised: this means that you do **not** have to write your name or your student number on the exam, but you are identified by the number we told you prior to the exam. Please check the correctness of the identifying number above before you start working!
- During the exam no written aids nor calculators or any other electronic device are allowed. **Phones must be switched off and stowed away** in your bag during the whole duration of the exam.
- A4-paper is provided. No other paper is allowed. Write with blue or black pens. **Do not use pencils, erasable pens, red or green ink, nor Tipp-Ex.**
- Write your solution to each exercise in the designated white space in the booklet. Leave enough ( $\approx 2$  cm) empty space on the margins (top, bottom and sides). If you need more space to write your solutions you can use the provided A4-paper, but please start every part on a new sheet of paper and **write your identifying number on every sheet of paper.**
- Please write neatly! Please do not put the graders in the unpleasant situation of being incapable of reading your solutions, as this will certainly not play in your favour!
- In part 3 and part 4 of this exam your answers do need to be properly justified. It is fine and allowed to use theorems/statements proved in class or in the homework (i. e. in Problem Sets 1–12) without reproving them (**unless otherwise stated**), but you should provide a precise statement of the result in question.
- Said  $0 \leq s \leq 60$  your total score on the exercises, your final grade in the exam will be *bounded from below* by  $\frac{s}{10}$ .
- The duration of the exam is **120 minutes**.

*Do not fill out this table!*

| part  | points | check |
|-------|--------|-------|
| 1     | [10]   |       |
| 2     | [10]   |       |
| 3     | [20]   |       |
| 4     | [20]   |       |
| total | [60]   |       |

grade:

**Part 1**

**[10 points]**

For each of the following 5 statement you have to establish whether it is true or false. Each correct answer is worth 2 points. An empty answer is worth 0 points. A wrong answer detracts 2 points (that is, it is worth  $-2$  points). If your total score in this exercise is negative, you will anyway be given 0 points (so, it is impossible to get a negative score on the whole exercise).

Insert your answers in the following grid. Write clearly **T** if the statement is true and **F** if the statement is false. We will accept also **R** if the statement is *richtig* (which is the German word for *true*).

Only the answers in the grid will be taken into consideration for grading.

|          |   |   |   |   |   |
|----------|---|---|---|---|---|
| Question | 1 | 2 | 3 | 4 | 5 |
| Answer   |   |   |   |   |   |

Let  $f : [0, \pi) \rightarrow \mathbb{R}$  be given by  $f(x) = x - 3 \cos(5x)$ . Let then  $g : (-\pi, \pi) \rightarrow \mathbb{R}$  denote the even extension of  $f$ , and let us consider its Fourier series

$$S_g(x) = \frac{a_0}{2} + \sum_{n \geq 1} (a_n \cos(nx) + b_n \sin(nx)).$$

1.  $a_0 > 0$

2.  $b_{90} < 0$

3.  $S_g(0) = -3/2$

4.  $a_5 = -\frac{1}{5\pi} - 3$

5.

$$\sum_{n \text{ odd}} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots = \frac{\pi^2}{8}$$

*Hint:* Compute  $S_g(\pi)$ .

**Part 2**

**[10 points]**

For each of the following 5 questions, you have to provide a *numeric or symbolic answer*. An example of a numeric answer is  $-27\sqrt{2}$ ; an example of a symbolic answer is  $a^3/b$ . Each correct answer is worth 2 points. An empty or wrong answer is worth 0 points.

Insert your answers in the following grid. Write clearly so that there is no ambiguity. Only the answers in the grid will be taken into consideration for grading. You do **not** have to provide any justification for your answers.

*Note: in the following questions, to be considered correct your answer needs to be explicit, i.e. it should not be given in the form of an unsolved integral or series.*

| Question | Answer |
|----------|--------|
| 1        |        |
| 2        |        |
| 3        |        |
| 4        |        |
| 5        |        |

Throughout this exercise, we let  $f(x) = e^{-x^2}$ .

*Note: We recall that the Fourier transform of  $h(x) = e^{-x^2/2}$  was computed in class and is given by  $\hat{h}(\xi) = \sqrt{2\pi}e^{-\xi^2/2}$ . In answering the following questions we advise you to exploit the properties of the Fourier transform whenever possible.*

1. What is the Fourier transform of  $f$ ?
2. What is the Fourier transform of  $g$ , where  $g(x) = f(x - 1)$ ?
3. What is the convolution product  $f * f$ ?
4. What is the value of  $\int_{\mathbb{R}} x^4 f(x) dx$ ?
5. What is the value  $u(x = 0, t = 1/200)$  where  $u$  is the (only integrable) solution to the heat equation

$$\begin{cases} u_t = 100u_{xx} & \text{for } x \in \mathbb{R}, t > 0, \\ u(x, 0) = f(x) & \text{for } x \in \mathbb{R}. \end{cases} \quad ?$$

**Part 3**

**[20 points]**

Consider the disc  $D := \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 4\}$ , and let  $u \in C^2(D; \mathbb{R})$  solve the elliptic boundary value problem

$$\begin{cases} \Delta u = 0 & \text{in } D \\ u = g & \text{in } \partial D \end{cases}$$

where  $g(x, y) = -1 + x^2 + 2y$ .

1. Write  $g$  in polar coordinates  $(r, \theta)$ ;
2. Determine  $u(0, 0)$ , i.e. the value of the solution in question at the center of the disc;
3. Determine  $\max_{(x,y) \in D} u(x, y)$  and  $\min_{(x,y) \in D} u(x, y)$  and, for each of them, the point(s) where they are attained;
4. Compute the (real) Fourier series expansion of  $g(2, \theta)$ ;
5. Determine the explicit expression of  $u$  as a series of functions.

*Hint:* Recall the trigonometric identity  $\cos(2\theta) = 2\cos^2(\theta) - 1$ .









**Part 4**

**[20 points]**

Consider the initial value problem

$$\begin{cases} u_{tt} = c^2 u_{xx} & \text{for } (x, t) \in \mathbb{R} \times \mathbb{R}_{\geq 0}, \\ u(x, 0) = \max \{4 - |x - 2|, 0\} & \text{for } x \in \mathbb{R}, \\ u_t(x, 0) = 0 & \text{for } x \in \mathbb{R}. \end{cases}$$

where  $c$  is a real, positive constant.

1. Determine  $u(x, t)$  for all  $(x, t) \in \mathbb{R} \times \mathbb{R}_{\geq 0}$ ;
2. Draw, as a function of  $c > 0$ , the graph of the value  $u(1, 1)$ ;
3. Determine the points  $(x, t) \in \mathbb{R} \times \mathbb{R}_{\geq 0}$  where the amplitude of the wave, i.e. the value of  $u$ , is maximum, and determine such a value;
4. Compute the value of  $\lim_{t \rightarrow +\infty} u(-3, t)$ ;
5. Suppose we now perform a modification of the initial datum  $u(x, 0)$  only for  $x > 0$  (while keeping it unchanged for  $x \leq 0$ ). What is the largest set of points  $(x, t) \in \mathbb{R} \times \mathbb{R}_{> 0}$  for which the value of  $u(x, t)$  may be affected by such a modification? Describe it by means of equations (in terms of  $c$ ), and draw it. Also: for what values of  $c$  will such modifications potentially affect  $u(-1, 3)$ ?





