

Mathematical Foundations for Finance

Exercise sheet 9

Please upload your solutions until Wednesday, 24/11/2021, 12:00 using the link on the course website.

Exercise 9.1 Let $(Y_k)_{k \in \mathbb{N}}$ be a sequence of independent random variables defined on a probability space (Ω, \mathcal{F}, P) and consider the filtration $\mathbb{F} = (\mathcal{F}_k)_{k \in \mathbb{N}_0}$ given by $\mathcal{F}_0 = \{\emptyset, \Omega\}$ and $\mathcal{F}_k = \sigma(Y_1, \dots, Y_k)$ for all $k \in \mathbb{N}$. Let $E[Y_k] = \mu$ and $\text{Var}(Y_k) = \sigma^2$ for all $k \in \mathbb{N}$ with $\mu \in \mathbb{R}$ and $\sigma^2 > 0$. Define additionally $X = (X_k)_{k \in \mathbb{N}_0}$ by

$$X_k = \sum_{j=1}^k Y_j \quad \text{for all } k \in \mathbb{N}_0.$$

- (a) Show that for any \mathbb{F} -adapted integrable process $Z = (Z_k)_{k \in \mathbb{N}_0}$, there exists a P -a.s. unique decomposition of Z into $Z = M + A$ with $M = (M_k)_{k \in \mathbb{N}_0}$ a (P, \mathbb{F}) -martingale and $A = (A_k)_{k \in \mathbb{N}_0}$ an \mathbb{F} -predictable integrable process with $A_0 = 0$.

Hint: This is the Doob decomposition. Show the existence by construction. You can start by rewriting $Z_k - Z_{k-1}$.

- (b) Using (a), explicitly derive the processes M and A in the Doob decomposition of X .

- (c) Explicitly derive the optional quadratic variation $[M] = ([M]_k)_{k \in \mathbb{N}_0}$ of the square-integrable martingale M from (b), and show that $M^2 - [M]$ is a martingale.

Hint: See Theorem V.1.1 in the lecture notes, and use that due to the condition $\Delta[M] = (\Delta M)^2$, we must have that $[M]_k - [M]_{k-1} = (M_k - M_{k-1})^2$.

- (d) Explicitly derive the predictable compensator $\langle M \rangle = (\langle M \rangle_k)_{k \in \mathbb{N}_0}$ of the process M from (b).

Hint: See the remark on page 79 in the lecture notes. Also use that if M is a square-integrable martingale, then $\langle M \rangle$ is integrable.

Exercise 9.2 A Poisson process with parameter $\lambda > 0$ with respect to a probability measure P and a filtration $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$ is a (real-valued) stochastic process $N = (N_t)_{t \geq 0}$ which is adapted to \mathbb{F} , $N_0 = 0$ P -a.s. and satisfies the following two properties:

- (PP1) For $0 \leq s < t$, the increment $N_t - N_s$ is independent (under P) of \mathcal{F}_s and is (under P) Poisson-distributed with parameter $\lambda(t - s)$, i.e.

$$P[N_t - N_s = k] = \frac{(\lambda(t - s))^k}{k!} e^{-\lambda(t - s)}, \quad k \in \mathbb{N}_0.$$

- (PP2) N is a counting process with jumps of size 1, i.e. for P -almost all $\omega \in \Omega$, the function $t \mapsto N_t(\omega)$ is right-continuous with left limits (RCLL), piecewise constant, \mathbb{N}_0 -valued, and increases by jumps of size 1.

Poisson processes form the cornerstone of jump processes, which are of importance in advanced financial modeling. Show that the following processes are (P, \mathbb{F}) -martingales:

- (a) $\tilde{N}_t := N_t - \lambda t$, $t \geq 0$. This process is also called a compensated Poisson process.

Hint: If $X \sim \text{Poi}(\lambda)$, then $E[X] = \lambda$.

(b) $\tilde{N}_t^2 - N_t$, $t \geq 0$, and $\tilde{N}_t^2 - \lambda t$, $t \geq 0$. Use these results to derive $[\tilde{N}]$ and $\langle \tilde{N} \rangle$.
Hint: If $X \sim \text{Poi}(\lambda)$, then $\text{Var}(X) = \lambda$.

(c) $S_t := e^{N_t \log(1+\sigma) - \lambda \sigma t}$, $t \geq 0$, where $\sigma > -1$. S is also called a *geometric Poisson process*.

Exercise 9.3 Let $(\Pi_n)_{n \in \mathbb{N}}$ be a sequence of refining partitions of $[a, b] \subseteq \mathbb{R}$ (in the sense that $\Pi_n \subseteq \Pi_{n+1}$ for all $n \in \mathbb{N}$) with $|\Pi_n| \rightarrow 0$ as $n \rightarrow \infty$. Let $p > 0$. We define for a function $f : \mathbb{R} \rightarrow \mathbb{R}$ its p -variation on $[a, b]$ along the sequence $(\Pi_n)_{n \in \mathbb{N}}$ as

$$V_p^{(a,b)}(f) := \lim_{n \rightarrow \infty} \sum_{t_i \in \Pi_n} |f(t_i) - f(t_{i-1})|^p,$$

assuming that the limit exists. Assume additionally that f is continuous on $[a, b]$.

(a) Show that if $V_{p^*}^{(a,b)}(f)$ is finite and non-zero for some $p^* > 0$, then $V_p^{(a,b)}(f) = \infty$ for all $p < p^*$.
Hint: Make sure to use the continuity of f . Use also that every function $f : \mathbb{R} \rightarrow \mathbb{R}$ that is continuous on a closed and bounded interval $[a, b]$ is also uniformly continuous on $[a, b]$.

(b) Show that if $V_{p^*}^{(a,b)}(f)$ is finite and non-zero for some $p^* > 0$, then $V_p^{(a,b)}(f) = 0$ for all $p > p^*$.