

### 7.1. (Non)homogeneous wave equation

(a) Let  $u = u(x, t)$  be a solution of the wave equation

$$\begin{cases} u_{tt} - c^2 u_{xx} = 1, & (x, t) \in \mathbb{R} \times (0, \infty), \\ u(x, 0) = 1, & x \in \mathbb{R}, \\ u_t(x, 0) = 1, & x \in \mathbb{R}, \end{cases}$$

Compute the explicit solution.

(b) Let  $u = u(x, t)$  be a solution of the wave equation

$$\begin{cases} u_{tt} - 2u_{xx} = 0, & (x, t) \in \mathbb{R} \times (0, \infty), \\ u(x, 0) = f(x), & x \in \mathbb{R}, \\ u_t(x, 0) = \sin(x), & x \in \mathbb{R}, \end{cases}$$

where  $f(x) = x$ , if  $|x| \leq 2$  and  $f(x) = 0$ , if  $|x| > 2$ . Is  $u$  smooth? Otherwise, where are the singularities of  $u$ ? Compute the explicit solution *after* answering these questions.

**7.2. Propagation of symmetries from initial data, II** Consider the general nonhomogeneous wave equation posed for  $-\infty < x < \infty$  and  $t > 0$ ,

$$\begin{cases} u_{tt} - c^2 u_{xx} = F(x, t), & (x, t) \in \mathbb{R} \times (0, \infty), \\ u(x, 0) = f(x), & x \in \mathbb{R}, \\ u_t(x, 0) = g(x), & x \in \mathbb{R}. \end{cases}$$

Take advantage of the uniqueness Theorem (4.4.6) in the notes to show that

(a) if  $f, g$  and  $F(\cdot, t)$  are odd/even functions, then  $u(\cdot, t)$  is itself odd/even.

(b) if  $f, g$  and  $F(\cdot, t)$  are periodic with same period  $T > 0$  (i.e.  $f(x + T) = f(x)$ ,  $g(x + T) = g(x)$  and  $F(x + T, t) = F(x, t)$  for all  $x \in \mathbb{R}$  and  $t > 0$ ), then  $u(\cdot, t)$  is itself periodic with period  $T$ .

**7.3. Wave equation on a ring** Let  $u : [0, 1] \times [0, \infty) \rightarrow \mathbb{R}$  be a solution of the wave equation

$$\begin{cases} u_{tt} - u_{xx} = 0, & (x, t) \in [0, 1] \times (0, \infty), \\ u(x, 0) = x - x^2, & x \in [0, 1], \\ u_t(x, 0) = 0, & x \in [0, 1], \\ u(0, t) = u(1, t), & t \in (0, \infty), \\ u_x(0, t) = u_x(1, t), & t \in (0, \infty). \end{cases}$$

Compute  $u(1/2, 2022)$ .

**7.4. Multiple choice** Cross the correct answer(s).

(a) Let  $u$  be solution of the homogeneous wave equation

$$\begin{cases} u_{tt} - 9u_{xx} = 0, & (x, t) \in \mathbb{R} \times (0, \infty), \\ u(x, 0) = f(x), & x \in \mathbb{R}, \\ u_t(x, 0) = g(x), & x \in \mathbb{R}. \end{cases}$$

Let  $h$  be a smooth function, and  $u_h$  be the solution of the above PDE with perturbed initial condition  $u_h(x, 0) = f(x)$  and  $(u_h)_t(x, 0) = g(x) + h(x)$ . Then,  $u(1, 2) = u_h(1, 2)$

- ☐ whenever  $h$  has compact support in  $[-5, 7]$ 
☐ only when  $h$  constantly equal to zero  
☐ whenever  $\int_{-5}^7 h(x) dx = 0$ 
☐ whenever  $h$  is equal to zero in  $[-5, 7]$

(b) Same question as (a), but when we perturb  $u_h(x, 0) = f(x) + h(x)$ ,  $(u_h)_t(x, 0) = g(x)$ .

- ☐ only when  $h$  constantly equal to zero
 ☐ always for  $h$  small enough  
☐ when  $h(x) = \sin(\pi(x + 1))$ 
☐ whenever  $h(-5) = h(7) = 0$

(c) Let  $u$  be solution of the homogeneous wave equation

$$\begin{cases} u_{tt} - u_{xx} = F(x), & (x, t) \in \mathbb{R} \times (0, \infty), \\ u(x, 0) = f(x), & x \in \mathbb{R}, \\ u_t(x, 0) = g(x), & x \in \mathbb{R}. \end{cases}$$

Suppose that  $F$ ,  $f$  and  $g$  are trigonometric polynomials as in Exercise 6.2 (b), with  $\int_0^{2\pi} g dx = 0$ . Then,  $u$  is

- ☐ never
 ☐ always for  $F \equiv 0$   
☐ always
 ☐ never unless  $F \neq f$

$2\pi$ -periodic in time<sup>1</sup>, that is  $u(x, t + 2\pi) = u(x, t)$  for all  $(x, t) \in \mathbb{R} \times (0, +\infty)$ .

---

<sup>1</sup>Be careful, this is not the same as being periodic in the  $x$  variable, as in Exercise 7.2.

## Extra exercises

**7.5. Strange wave equation** Show that the following partial differential equation admits a solution

$$\begin{cases} u_{tt} - u_{xx} &= \frac{u_t^2 - u_x^2}{2u}, & (x, t) \in \mathbb{R} \times (0, \infty), \\ u(x, 0) &= x^4, & x \in \mathbb{R}, \\ u_t(x, 0) &= 0, & x \in \mathbb{R}. \end{cases}$$

*Hint: Consider the function  $v(x, t) = \sqrt{u(x, t)}$ . What equation does it satisfy?*