

EQUIVALENCE OF SIMPLICIAL & SINGULAR HOMOLOGY

Recall that simplicial homology was defined in terms of a Δ -complex decomposition of X , via a collection of maps

$$\delta_\alpha: \Delta^p \rightarrow X.$$

We then defined the chains to be the free abelian groups on the p -simplices, $\Delta_p(X)$. We will now show that if a Δ -complex structure is chosen, then its simplicial homology coincides with the singular homology of the space X . We will do this by induction on the skeleton $X^{(k)}$ consisting of all simplices of dimension k or less, and so we would like to use a relative version of simplicial homology.

LEMMA 1

For a wedge sum $\bigvee_{\alpha} X_{\alpha}$, the inclusions $i_{\alpha}: X_{\alpha} \hookrightarrow \bigvee_{\alpha} X_{\alpha}$ induce an isomorphism

$$\bigoplus_{\alpha} i_{\alpha*} : \bigoplus_{\alpha} \tilde{H}_p(X_{\alpha}) \rightarrow \tilde{H}_p\left(\bigvee_{\alpha} X_{\alpha}\right),$$

provided that the wedge sum is formed at basepoints $x_{\alpha} \in X_{\alpha}$ such that the pairs (X_{α}, x_{α}) are good.

Recall

The wedge sum is a one-point union of a family of topological spaces:

$$\bigvee_{\alpha} (X_{\alpha}, x_{\alpha}) = \left(\bigsqcup_{\alpha} X_{\alpha} \right) / \left\{ x_{\alpha} \right\}_{\alpha}$$

quotient of the disjoint union of the X_{α} 's by the equivalence relation that identifies all x_{α} 's with each other and makes no other identifications

Proof

Since (X_α, x_α) 's are good pairs,

$(\bigsqcup_\alpha X_\alpha, \{x_\alpha : \alpha \in I\})$ is a good pair,

so

$$H_p \left(\bigsqcup_\alpha X_\alpha, \{x_\alpha : \alpha \in I\} \right) \xrightarrow[\cong]{\cong} H_p \left(\bigsqcup_\alpha X_\alpha / \{x_\alpha\}_\alpha \right) \\ \parallel \\ \tilde{H}_p \left(\bigvee_\alpha X_\alpha \right)$$

Furthermore, for pairs of topological spaces we have

$$H_p \left(\bigsqcup_\alpha (A_\alpha, B_\alpha) \right) \cong \bigoplus_\alpha H_p (A_\alpha, B_\alpha).$$

↙ induced by inclusions

It follows from here that

$$\tilde{H}_p \left(\bigvee_\alpha X_\alpha \right) \cong \bigoplus_\alpha H_p (X_\alpha, x_\alpha)$$

We did an example a while back

$$\rightarrow \cong \bigoplus_\alpha \tilde{H}_p (X_\alpha)$$


and showed that

$$H_p(X, x_0) \cong \tilde{H}_p(X).$$

EXAMPLE

$$\text{Let } X = \underbrace{S^n \vee S^n \vee \dots \vee S^n}_{\text{a finite wedge of } h \text{ } n\text{-spheres}}$$

a finite wedge
of h n -spheres



then by Lemma 1

$$\tilde{H}_p(\vee S^n) \cong \tilde{H}_p(S^n) \oplus \dots \oplus \tilde{H}_p(S^n)$$

$$= \begin{cases} \oplus_h \mathbb{Z} & p = n \\ 0 & \text{otherwise} \end{cases}$$