

DEFINITION (RELATIVE SIMPLICIAL HOMOLOGY)

Let X be a Δ -complex & A a Δ -subcomplex.

We define the quotient chain complex

$$\Delta_p(X, A) = \frac{\Delta_p(X)}{\Delta_p(A)}$$

relative simplicial chains

with ∂ restrictions of simplicial boundary maps. The homology of this chain complex is denoted by $H_p^\Delta(X, A)$.

THEOREM

maps that give the Δ -complex structure
the characteristic map of any p -simplex in a Δ -complex decomposition

of X may be viewed as a singular p -simplex, hence we have the inclusion chain map i_c

$$\begin{array}{ccccccc} \cdots & \rightarrow & \Delta_p(X, A) & \rightarrow & \Delta_{p-1}(X, A) & \rightarrow & \Delta_{p-2}(X, A) \rightarrow \cdots \\ & & \downarrow i_c & & \downarrow i_c & & \downarrow i_c \\ \cdots & \rightarrow & S_p(X, A) & \rightarrow & S_{p-1}(X, A) & \rightarrow & S_{p-2}(X, A) \rightarrow \cdots \end{array}$$

The induced homomorphism

$H_p^\Delta(X, A) \rightarrow H_p(X, A)$ is an isomorphism.

Taking $A = \phi$, we obtain the equivalence of absolute singular and simplicial homology.

Proof

(i) special case : $A = \phi$, X is finite-dimensional
We denote by $X^{(k)}$ the k -skeleton of X .

Then we have the following LES

$$\begin{array}{ccccccccc} \rightarrow H_{p+1}^\Delta(X^{(k)}, X^{(k-1)}) & \rightarrow & H_p^\Delta(X^{(k-1)}) & \rightarrow & H_p^\Delta(X^{(k)}) & \rightarrow & H_p^\Delta(X^{(k)}, X^{(k-1)}) & \rightarrow & H_p^\Delta(X^{(k-1)}) \\ \downarrow \textcircled{1} & & \downarrow \textcircled{2} & & \downarrow \textcircled{3} & & \downarrow \textcircled{4} & & \downarrow \textcircled{5} \\ \rightarrow H_{p+1}(X^{(k)}, X^{(k-1)}) & \rightarrow & H_p(X^{(k-1)}) & \rightarrow & H_p(X^{(k)}) & \rightarrow & H_p(X^{(k)}, X^{(k-1)}) & \rightarrow & H_p(X^{(k-1)}) \end{array}$$

First we show that $\textcircled{1}$ & $\textcircled{4}$ are an isomorphism.

Note that $\Delta_p(X^{(k)}, X^{(k-1)}) = 0$ for $p \neq k$

& furthermore, $\Delta_k(X^{(k)}, X^{(k-1)})$ is a free abelian group generated by k -simplices in X

$$\text{Hence, } H_p^\Delta(X^{(k)}, X^{(k-1)}) = \begin{cases} 0 & p \neq k \\ \bigoplus_{\substack{\# k-SX \\ \text{in } X}} \mathbb{Z} & p = k \end{cases}$$

To compute $H_p(X^{(k)}, X^{(k-1)})$ note that $X^{(k-1)}$ is a strong deformation retract of its neighborhood in $X^{(k)}$.

(proposition A.5. on page 523 in Hatcher).

So

$$H_p(X^{(k)}, X^{(k-1)}) \xrightarrow{\cong} \tilde{H}_p\left(\frac{X^{(k)}}{X^{(k-1)}}\right)$$

$$\frac{X^{(k)}}{X^{(k-1)}} = \bigvee_{\substack{\# k-SX \\ \text{in } X}} S^k \Rightarrow$$

$$\tilde{H}_p\left(\frac{X^{(k)}}{X^{(k-1)}}\right) = \tilde{H}_p\left(\bigvee_{\substack{\# k-SX \\ \text{in } X}} S^k\right) = \begin{cases} 0 & p \neq k \\ \bigoplus_{\substack{\# k-SX \\ \text{in } X}} \mathbb{Z} & p = k \end{cases}$$

$\Rightarrow H_p^\Delta(X^{(k)}, X^{(k-1)}) \rightarrow H_p(X^{(k)}, X^{(k-1)})$ is an isomorphism.

$$\begin{array}{ccccccc}
 \rightarrow H_{p+1}^\Delta(X^{(k)}, X^{(k-1)}) & \rightarrow & H_p^\Delta(X^{(k-1)}) & \rightarrow & H_p^\Delta(X^{(k)}) & \rightarrow & H_p^\Delta(X^{(k)}, X^{(k-1)}) \rightarrow H_p^\Delta(X^{(k-1)}) \\
 \text{?} \parallel \downarrow \textcircled{1} & & \downarrow \textcircled{2} & & \downarrow \textcircled{3} & & \text{?} \parallel \downarrow \textcircled{4} & & \downarrow \textcircled{5} \\
 \rightarrow H_{p+1}(X^{(k)}, X^{(k-1)}) & \rightarrow & H_p(X^{(k-1)}) & \rightarrow & H_p(X^{(k)}) & \rightarrow & H_p(X^{(k)}, X^{(k-1)}) \rightarrow H_p(X^{(k-1)})
 \end{array}$$

So $\textcircled{1}$ & $\textcircled{4}$ are isomorphisms.

All homology groups of the empty set are trivial. Assume that

$H_p^\Delta(X^{(j)}) \rightarrow H_p(X^{(j)})$ are isomorphisms for $j \leq k-1$ and for all p .

By induction on k we may assume that the second $\textcircled{2}$ and the fifth $\textcircled{5}$ map are isomorphisms as well.

Then $\textcircled{3}$ is an isomorphism by the five-lemma.

(ii) X infinite dimensional, $A = \emptyset$

FACT A compact set in X can meet X in only finitely many open simplices of X , i.e. simplices

with their proper faces deleted.

Proof of the fact

If a compact set C intersected infinitely many open simplices, it would contain an infinite sequence of points x_i each lying in a different open simplex. Then

$$U_i = X - \bigcup_{j \neq i} \{x_j\}$$

are open since their preimages under the characteristic maps of all simplices are open. They also form an open cover with no subcover.

Let us show that

① $H_p^\Delta(X) \rightarrow H_p(X)$ is surjective

Let $m \in H_p(X)$, $m = [c]$, $c \in S_p(X)$, $\partial c = 0$.

c is a finite linear combination of singular simplices, whose images are compact subsets of X . So the images of simplices are a compact set. Using the fact above, we deduce that these images are in $X^{(k)}$ for k big enough. $\Rightarrow c \in S_p(X^{(k)})$.

By (i) we know that $H_p^\Delta(X^{(k)}) \rightarrow H_p(X^{(k)})$ is an isomorphism, so $v \in \Delta_p(X^{(k)})$

exists, $\partial v = 0$ such that $[v]$

gets mapped to $[c] \in H_p(X^{(k)})$.

Now $v \in \Delta_p(X^{(k)}) \subset \Delta_p(X)$ &

$[v]$ is mapped to $[c] \in H_p(X)$.

$$\begin{array}{ccc}
 H_p^\Delta(X^{(k)}) & \longrightarrow & H_p(X^{(k)}) \\
 \downarrow & & \downarrow \\
 H_p^\Delta(X) & \longrightarrow & H_p(X)
 \end{array}$$

② $H_p^\Delta(X) \rightarrow H_p(X)$ is injective

Let $c \in \Delta_p(X)$, $[c]$ is in the kernel of the above homomorphism. This means that c is a boundary of some singular $(p+1)$ -chain $b \in S_{p+1}(X)$.

Both chains, c and b , lie in $X^{(k)}$ for a large enough k (the argument is the same as before). This means that c represents an element in the kernel of

$$\begin{array}{ccc}
 H_p^\Delta(X^{(k)}) & \xrightarrow{\cong} & H_p(X^{(k)}) \\
 [c] & \longmapsto & 0
 \end{array} \quad (i)$$

$$\Rightarrow [c] = 0 \in H_p^\Delta(X^{(k)}) \Rightarrow [c] = 0$$

in $H_p^\Delta(k)$.

(iii) X, A general, $A \subset X$. Then LES for (X, A) yields

$$\begin{array}{ccccccccc} \rightarrow H_{p+1}^\Delta(X, A) & \rightarrow & H_p^\Delta(A) & \rightarrow & H_p^\Delta(X) & \rightarrow & H_p^\Delta(X, A) & \rightarrow & H_p^\Delta(A) \rightarrow \\ & & \downarrow \cong (ii) & & \downarrow \cong (ii) & & \downarrow & & \downarrow \cong (i) \\ \rightarrow H_{p+1}(X, A) & \rightarrow & H_p(A) & \rightarrow & H_p(X) & \rightarrow & H_p(X, A) & \rightarrow & H_p(A) \rightarrow \end{array}$$

By the 5-lemma $H_p^\Delta(X, A) \rightarrow H_p(X, A)$

is also an isomorphism.