

# SINGULAR HOMOLOGY

## Definition

A SINGULAR  $n$ -simplex in  $X$  is a map  $\sigma: \Delta^n \rightarrow X$ .

The word singular is used to express the idea that  $\sigma$  need not be a nice embedding, but can have 'singularities' where its image does not look like a simplex. All that is required is that  $\sigma$  is continuous.

## Definitions

$S_n(X)$  is the free abelian group generated by all the singular  $n$ -simplices  $\sigma: \Delta^n \rightarrow X$  of  $X$ .

We call  $S_n(X)$  the GROUP OF SINGULAR  $n$ -CHAINS of  $X$ .

A singular  $n$ -chain is a (finite) formal sum

$$\sum_{\sigma: \Delta^n \rightarrow X} n_\sigma \cdot \sigma, \quad n_\sigma \in \mathbb{Z}.$$

The **BOUNDARY MAP** is defined by the same formula as before:

$$\partial_n(\sigma) = \sum_i (-1)^i \sigma \Big|_{[v_0, \dots, \hat{v}_i, \dots, v_n]}$$

$\sigma \Big|_{[v_0, \dots, \hat{v}_i, \dots, v_n]}$  is regarded as a map  $\Delta^{n-1} \rightarrow X$  via the canonical identification of  $[v_0, \dots, \hat{v}_i, \dots, v_n]$

with  $\Delta^{n-1}$ , preserving the ordering of the vertices

the proof that  $\partial_n \circ \partial_{n+1} = 0$  works the same as in the simplicial homology case.

**Notation:** We often denote all  $\partial_n$  by  $\partial$  and write  $S_n(x) \xrightarrow{\partial} S_{n-1}(x)$  &  $\partial \circ \partial = 0$ .

**Definition**

$C_n(x) = (S_n(x), \partial_n)_n$  is a chain complex.

The **SINGULAR HOMOLOGY** groups  $\downarrow$  cycles  
are

$$H_n(x) = \ker \partial_n$$

$$\ker \partial_n = Z_n(x)$$

$$\operatorname{Im} \partial_{n+1} = B_n(x)$$

$$\operatorname{Im} \partial_{n+1}$$

boundaries  $\uparrow$

**Example**

$X$  point. What are the homology groups of  $X$ ?

For each dimension  $n \geq 0$  we have exactly one singular simplex

$\mathcal{G}_n: \Delta^n \rightarrow X$ , so,  $S_n(X) = \mathbb{Z} \cdot \mathcal{G}_n$ .

We now calculate  $\partial_n: S_n(X) \rightarrow S_{n-1}(X)$ .

$\partial_n(\mathcal{G}_n)$  = an alternating sum of  $(n+1)$  elements each of which is  $\mathcal{G}_{n-1}$

$$\partial_n(\mathcal{G}_n) = \begin{cases} 0 & n = \text{odd} \\ \mathcal{G}_{n-1} & n \text{ is even } > 0 \\ 0 & n = 0 \end{cases}$$

$$\begin{array}{ccccccc} \dots & \rightarrow & S_3(X) & \xrightarrow{\partial_3} & S_2(X) & \xrightarrow{\partial_2} & S_1(X) & \xrightarrow{\partial_1} & S_0(X) & \rightarrow & 0 \\ & & \parallel \mathbb{Z} & & \parallel \mathbb{Z} & \cong & \parallel \mathbb{Z} & \xrightarrow{0} & \parallel \mathbb{Z} & \rightarrow & 0 \\ \dots & \rightarrow & \mathbb{Z} & \xrightarrow{0} & \mathbb{Z} & \xrightarrow{\cong} & \mathbb{Z} & \xrightarrow{0} & \mathbb{Z} & \rightarrow & 0 \end{array}$$

$\partial_n$  is an isomorphism for even  $n > 0$  and the zero map when  $n$  is odd.

Cycles:

$$Z_n(x) = \begin{cases} \mathbb{Z} & n = \text{odd} \\ 0 & n = \text{even} \ \& \ n > 0 \\ \mathbb{Z} & n = 0 \end{cases}$$

Boundaries

$$B_n(x) = \begin{cases} \mathbb{Z} & n = \text{odd} \\ 0 & n = \text{even} \ \& \ n > 0 \\ 0 & n = 0 \end{cases}$$

$$H_n(x) = \begin{cases} 0 & n = \text{odd} \\ 0 & n = \text{even} \ \& \ n > 0 \\ \mathbb{Z} & n = 0 \end{cases}$$

$$= \begin{cases} \mathbb{Z} & n = 0 \\ 0 & n \neq 0 \end{cases}$$

## FUNCTORIAL PROPERTIES

Let  $f: X \rightarrow Y$  be a map between the spaces  $X$  &  $Y$ . For every