

Cycles:

$$Z_n(x) = \begin{cases} \mathbb{Z} & n = \text{odd} \\ 0 & n = \text{even} \ \& \ n > 0 \\ \mathbb{Z} & n = 0 \end{cases}$$

Boundaries

$$B_n(x) = \begin{cases} \mathbb{Z} & n = \text{odd} \\ 0 & n = \text{even} \ \& \ n > 0 \\ 0 & n = 0 \end{cases}$$

$$H_n(x) = \begin{cases} 0 & n = \text{odd} \\ 0 & n = \text{even} \ \& \ n > 0 \\ \mathbb{Z} & n = 0 \end{cases}$$

$$= \begin{cases} \mathbb{Z} & n = 0 \\ 0 & n \neq 0 \end{cases}$$

## FUNCTORIAL PROPERTIES

Let  $f: X \rightarrow Y$  be a map between the spaces  $X$  &  $Y$ . For every

Singular  $n$ -simplex  $\sigma: \Delta^n \rightarrow X$ ,  
 we get a new singular simplex  
 induced by  $f \circ \sigma: \Delta^n \rightarrow Y$ .

Extending linearly we get  
 a homomorphism  $f_c: C_n(X) \rightarrow C_n(Y)$   
 defined by

$$f_n = S_n(f) : S_n(X) \rightarrow S_n(Y)$$

$$f_m \left( \sum_{\sigma} n_{\sigma} \cdot \sigma \right) = \sum_{\sigma} n_{\sigma} (f \circ \sigma)$$

Proposition

$$f_{n-1} \circ \partial_n = \partial_n \circ f_n \quad (f_c \circ \partial = \partial \circ f_c)$$

Proof

$$\begin{aligned} f_{n-1} \circ \partial_n(\sigma) &= f_{n-1} \left( \sum_i (-1)^i \sigma|_{[v_0, \dots, \hat{v}_i, \dots, v_n]} \right) \\ &= \sum_i (-1)^i f \circ \sigma|_{[v_0, \dots, \hat{v}_i, \dots, v_n]} \end{aligned}$$

$$= \partial_n (f \circ \sigma) = \partial_n \circ f_n (\sigma)$$



thus we have a commutative diagram

$$\begin{array}{ccccccc}
 \cdots & \rightarrow & C_{n+1}(X) & \xrightarrow{\partial_{n+1}} & C_n(X) & \xrightarrow{\partial_n} & C_{n-1}(X) & \rightarrow \cdots \\
 & & \downarrow f_{n+1} & & \downarrow f_n & & \downarrow f_{n-1} & \\
 \cdots & \rightarrow & C_{n+1}(Y) & \xrightarrow{\partial_{n+1}} & C_n(Y) & \xrightarrow{\partial_n} & C_{n-1}(Y) & \rightarrow \cdots
 \end{array}$$

Homomorphisms  $f_c = C_n(X) \rightarrow C_n(Y)$  that satisfy  $f_c \circ \partial = \partial \circ f_c$  are called **CHAIN MAPS** from the singular chain complex of  $X$  to that of  $Y$ .

## COROLLARY

$$\textcircled{1} f_c(Z_n(x)) \subset Z_n(Y)$$

$$f_c(B_n(x)) \subset B_n(Y)$$

In particular  $f$  induces, via  $f_c$ ,  
a homomorphism  $f_*: H_n(x) \rightarrow H_n(Y)$ ,  
by  $f_*([c]) = [f_c(c)]$ .

$$\textcircled{2} x \xrightarrow{f} y \xrightarrow{g} z \Rightarrow$$

$$(g \circ f)_* = g_* \circ f_*: H_n(x) \rightarrow H_n(z)$$

$$\& (\text{id}_x)_* = \text{id}_{H_n(x)}$$

Proof

$$\textcircled{1} \text{ if } c \in Z_n(x) \text{ (ie. } \partial c = 0) \Rightarrow$$

$$\partial f_c(c) = f_c(\underbrace{\partial c}_0) = 0$$

$$\Rightarrow f_c(c) \in Z_W(Y).$$

$$\text{If } c = \alpha d, d \in S_{n+1}(x) \Rightarrow$$

$$f_c(c) = f_c(\alpha d) = \alpha f_c(d) \in B_n(Y).$$

$\Rightarrow f$  induces a homomorphism

$$\underbrace{\frac{Z_n(x)}{B_n(x)}}_{H_n(x)} \xrightarrow{f_*} \underbrace{\frac{Z_n(Y)}{B_n(Y)}}_{H_n(Y)}$$

② Exercise.

Notation:

$$f_*, H(f): H_n(x) \rightarrow H_n(Y)$$

$f_*$  is called the map induced by

$f$  in homology.

### COROLLARY

If  $f: X \rightarrow Y$  is a homeomorphism, then  $f_*: H_n(X) \rightarrow H_n(Y)$  is an isomorphism  $\forall n$ .

Proof

Put  $g := f^{-1}: Y \rightarrow X$ , so  $f \circ g = \text{id}_Y$ ,

$$g \circ f = \text{id}_X.$$

$$\text{id}_{H_n(Y)} = (\text{id}_Y)_* = (f \circ g)_* =$$

$$= f_* \circ g_*: H_n(Y) \rightarrow H_n(Y)$$

$$\text{id}_{H_n(X)} = (\text{id}_X)_* = (g \circ f)_* =$$

$$= g_x \circ f_x : H_n(X) \rightarrow H_n(X).$$

## THE ZEROTH HOMOLOGY GROUP (Bredon)

$X$  space. What is  $H_0(X)$ ?

A 0-simplex  $\sigma: \Delta^0 \rightarrow X$  is just  
 $\parallel$   
 point

a choice of a point in  $X$ .

A 0-chain in  $X$  is a finite formal sum  $c = \sum_{x \in X} n_x \cdot x$ . Clearly,  $\partial(c) = 0$ .

Define  $\varepsilon(c) = \sum_{x \in X} n_x \in \mathbb{Z}$ .

Easy to check  $\varepsilon: S_0(X) \rightarrow \mathbb{Z}$  is  
 a homomorphism.

Let  $\sigma$  be a singular 1-simplex,

$\partial : \Delta^1 \rightarrow X$ , Put  $x_0 = \partial(0)$ ,  $x_1 = \partial(1)$ .

$$\partial \partial = \partial(1) - \partial(0) = x_1 - x_0$$

$$\Rightarrow \varepsilon(\partial \partial) = 1 - 1 = 0.$$

So for each 1-dimensional chain  $d$  we have  $\varepsilon(\partial d) = 0 \Rightarrow \varepsilon(B_0(x)) = 0$ .

It follows that  $\varepsilon$  induces a homomorphism

$$\varepsilon_x : H_0(x) \rightarrow \mathbb{Z}.$$

Both  $\varepsilon$  and  $\varepsilon_x$  are called

**AUGMENTATION.**



## Theorem

If  $X \neq \emptyset$  is path-connected,  
then  $\varepsilon_x : H_0(X) \rightarrow \mathbb{Z}$  is an  
isomorphism. Moreover,  $H_0(X) \cong \mathbb{Z} \cdot [x]$ ,  
where  $x \in X$  is any point.

## Proof


Clearly,  $\varepsilon_x([y]) = 1 \quad \forall y \in X$ .

So  $\varepsilon_x$  is surjective.

We must show that  $\varepsilon_x$  is injective.

Fix  $x_0 \in X$ .  $\forall x \in X$  choose a path

$\gamma_x : I \rightarrow X$  with  $\gamma_x(0) = x_0$ ,  $\gamma_x(1) = x$ .



If we view  $m_x$  as a singular  
1-simplex,  $\partial m_x = x - x_0 \in S_0(X)$ .

Let  $c \in S_0(X)$ ,  $c = \sum_x n_x \cdot x$ , be a  
0-chain. Assume  $[c] \in \ker(\mathcal{E}_*)$ ,

$$\text{we } \sum_x n_x = 0.$$

Consider the 1-chain  $\sum n_x m_x \in S_1(X)$ .

$$\partial(\sum n_x m_x) = \sum n_x (x - x_0) = \sum n_x \cdot x -$$

$$\underbrace{\left(\sum n_x\right)}_{=0} \cdot x_0 = c \Rightarrow c \in B_0(X).$$

$$\Rightarrow [c] = 0 \Rightarrow \ker \mathcal{E}_* = 0.$$

Proposition

Let  $X \neq \emptyset$  and denote by  $\mathcal{C}$  the  
set of path-connected components

of  $X$ .  $\forall \alpha \in \mathcal{C}$ , denote by  $X_\alpha \subset X$   
the path-connected component  
corresponding to  $\alpha$ . Then

$$H_n(X) \cong \bigoplus_{\alpha \in \mathcal{C}} H_n(X_\alpha), \text{ where}$$

this isomorphism is induced by  
inclusions. In particular,

$$H_0(X) \cong \bigoplus_{\alpha \in \mathcal{C}} H_0(X_\alpha)$$

$$\cong \bigoplus_{\alpha \in \mathcal{C}} \mathbb{Z}$$

Proof

Since a singular simplex always  
has a path-connected image,

$S_n(x)$  splits as the direct sum of

its subgroups  $S_n(x_\alpha)$ . The boundary maps  $\partial_n$  preserve this direct sum decomposition, taking  $S_n(x_\alpha)$  to  $S_{n-1}(x_\alpha)$ , so  $\ker \partial_n$  and  $\text{Im} \partial_{n+1}$  split similarly as direct sums. Hence, homology groups also split,

$$H_n(X) \cong \bigoplus_{\alpha \in \mathcal{E}} H_n(x_\alpha).$$

## REDUCED HOMOLOGY GROUPS

(Hatcher)

It is often convenient to have a slightly modified version of homology for which a point has trivial homology in all dimensions, including 0. This

is done by defining **REDUCED**  
**HOMOLOGY GROUPS**  $\tilde{H}_n(X)$  to  
 be the homology groups of  
 the **AUGMENTED CHAIN**  
**COMPLEX**

$$\dots \rightarrow S_2(X) \xrightarrow{\partial_2} S_1(X) \xrightarrow{\partial_1} S_0(X) \xrightarrow{\epsilon} \mathbb{Z} \rightarrow 0$$

where  $\epsilon$  is the augmentation.  
 Usually, we require  $X$  to be  
 nonempty to avoid a non-trivial  
 homology group in  $\dim -1$ .

Then  $\tilde{H}_0(X)$  is the kernel of  
 $\epsilon^*$  so that  $H_0(X) \cong \tilde{H}_0(X) \oplus \mathbb{Z}$ .