

Solutions 7

1. Characterization of the Sphere

Prove the following lemma due to H. Hopf:

Lemma. Let M be a compact, connected, m -dimensional submanifold of \mathbb{R}^{m+1} . Suppose that for each vector $v \in S^m$ there exists $\lambda = \lambda(v) \in \mathbb{R}$ such that M is symmetric with respect to reflections on the hyperplane $E_{v,\lambda} := \{x \in \mathbb{R}^{m+1} : \langle x, v \rangle = \lambda\}$, then M is a sphere.

Solution. a) First of all, notice that $\langle x, v \rangle = \lambda(v)$ if and only if $\langle x - \lambda(v)v, v \rangle = 0$, hence $E_{v,\lambda(v)} = T_{\lambda(v)v}(E_{v,0})$ and in fact $E_{v,\lambda} = T_{\lambda v}(E_{v,0})$ for all λ , where $T_{\lambda v}: x \mapsto x + \lambda v$. Moreover the reflection on the hyperplane $E_{v,0}$ is given by $R_{v,0}: z \mapsto z - 2\langle z, v \rangle v$. Hence the reflection on the hyperplane $E_{v,\lambda(v)}$ is given by

$$R_{v,\lambda(v)} = T_{\lambda(v)v} \circ R_{v,0} \circ T_{-\lambda(v)v} \quad z \mapsto z - 2\langle v, z - \lambda(v)v \rangle v.$$

Up to translating M , we might assume that M is symmetric with respect to the reflections on the hyperplanes $E_i := E_{e_i,0}$ (which is given by changing the sign of the i -th coordinate)¹. By applying successively the reflections on the hyperplanes E_1, \dots, E_{m+1} we obtain that M is preserved by the map $x \mapsto -x$.

This implies that $\lambda(v) = 0$ for all $v \in S^m$. Indeed, first notice that if M is symmetric with respect to $E_{v,\lambda}$, then M is symmetric with respect to $E_{-v,\lambda}$. This follows because M is preserved by the map $x \mapsto -x$ and also $R_{-v,\lambda}(z) = -R_{v,\lambda}(-z)$. If $\lambda(v) \neq 0$ we can use subsequent reflections on the parallel hyperplanes $E_{v,\lambda(v)} \neq E_{-v,\lambda(v)}$ to produce an unbounded sequence of points in M . This is not possible by compactness, hence $\lambda(v) = 0$.

Now let $p \in M \setminus \{0\}$. For every point $q \in S_{|p|}(0)$, the sphere with radius $|p|$ around 0, there exists a reflection $R_{v,0}$ on the hyperplane $E_{v,0}$ (explicitly $v := \frac{q-p}{|q-p|}$) sending p to q , so $q \in M$. Hence $S_{|p|}(0)$ is contained in M . If M contains any other point p' with $|p'| \neq |p|$, then the same argument shows that M contains also the sphere of radius $|p'|$ and by connectedness also the region between the two spheres, contradicting the m -dimensionality of M .

¹Denote by $T_1: \mathbb{R}^{m+1} \rightarrow \mathbb{R}^{m+1}$ the translation by $-\lambda(e_1) \cdot e_1$ and show that $T_1(M)$ is symmetric with respect to reflections on the hyperplane $E_{e_1,0}$. Notice that $T_1(M)$ is still symmetric with respect to the hyperplanes $E_{e_i,\lambda(e_i)}$ for $i = 2, \dots, m+1$. Repeat for T_2, \dots, T_{m+1} .

2. Non-positively Curved Surfaces

Let $M \subset \mathbb{R}^3$ be a surface with Gauss curvature $K \leq 0$. Prove the following assertions (we assume $a < b$).

- a) There is no simple geodesic loop (in particular no simple C^∞ -closed geodesic) $c: [a, b] \rightarrow M$ whose trace bounds a topological disk in M .
- b) There is no pair of injective geodesics $c_1, c_2: [a, b] \rightarrow M$ such that $c_1(a) = c_2(a)$ and $c_1(b) = c_2(b)$ are the only common points and the union of the traces bounds a topological disk.
- c) If M is homeomorphic to a cylinder and $K < 0$, then there is no pair of simple C^∞ -closed geodesics $c_1, c_2: [a, b] \rightarrow M$ with different traces.

Solution. a) Suppose that there is a geodesic c bounding a simply connected region D homeomorphic to a disc. Denote by α the external angle in $c(a) = c(b)$. By the Gauss-Bonnet Theorem (Theorem 6.3) we have

$$\int_D K \, dA + \alpha = 2\pi,$$

which is not possible as $K \leq 0$ and $\alpha \in [-\pi, \pi]$.

b) Suppose that the geodesics c_1 and c_2 enclose a compact simply connected region D homeomorphic to a disc. Denote by α_1, α_2 the external angles between them in p and in q , respectively. Since c_1 and c_2 are geodesics, it holds $\kappa_g = 0$. Hence by the Gauss-Bonnet Theorem (Theorem 6.3) we have

$$\int_D K \, dA + \alpha_1 + \alpha_2 = 2\pi.$$

Since $K \leq 0$ it follows that $\alpha_1 + \alpha_2 \geq 2\pi$. Moreover, since $\alpha_1, \alpha_2 \in [-\pi, \pi]$ it must hold $\alpha_1 = \alpha_2 = \pi$. Therefore $\dot{c}_1(0) = \dot{c}_2(0)$ and thus by uniqueness of geodesics $c_1 = c_2$, a contradiction.

c) Suppose that there are two geodesics c_1 and c_2 . From a) it follows that c_1 and c_2 cannot enclose a disc and cannot intersect (else, they would enclose discs and hence coincide). Therefore they must enclose an annulus.

We parametrize c_1, c_2 such that their orientation coincides with the one of the domain that they enclose. Choose two points p on c_1 and q on c_2 and a curve c connecting p to q , which doesn't intersect c_1 and c_2 in any other point. Then the concatenation $c_1 \cup c \cup c_2 \cup -c$ encloses a compact simply connected region D homeomorphic to a disc. It holds

$$\int_{c_1 \cup c \cup c_2 \cup -c} \kappa_g = \int_{c_1} \kappa_g + \int_c \kappa_g + \int_{c_2} \kappa_g + \int_{-c} \kappa_g = \int_c \kappa_g - \int_c \kappa_g = 0.$$

Since the external angles sum to 2π , the Gauss-Bonnet Theorem gives

$$0 > \int_D K \, dA = 2\pi - \sum_{i=1}^4 \alpha_i = 0,$$

which gives the desired contradiction.

3. Gauss Map of the Torus

- a) Let $f: U \rightarrow \mathbb{R}^{m+1}$, $U \subset \mathbb{R}^m$ open, be an immersion with Gauss map $\nu: U \rightarrow S^m$. Assuming that ν is an immersion, prove that

$$A(\nu) = \int_U |K| \sqrt{\det(g_{ij})} \, dx.$$

- b) Let $T \subset \mathbb{R}^3$ be a torus. Describe the image of the Gauss map and prove that

$$\int_T K \, dA = 0,$$

without using the Theorem of Gauss-Bonnet.

Solution. a) From Weingarten's equation (Lemma 4.8(2)) it follows that

$$\nu_i = - \sum_{k=1}^m h^k_i f_k$$

and hence

$$\begin{aligned} \langle \nu_i, \nu_j \rangle &= \left\langle - \sum_{k=1}^m h^k_i f_k, - \sum_{l=1}^m h^l_j f_l \right\rangle \\ &= \sum_{k=1}^m \sum_{l=1}^m h^k_i h^l_j \langle f_k, f_l \rangle \\ &= \sum_{k=1}^m \sum_{l=1}^m h^k_i h^l_j g_{kl} \\ &= \sum_{k=1}^m \sum_{l=1}^m h^i_k g_{kl} h^l_j \end{aligned}$$

So we get

$$\begin{aligned} \det(\langle \nu_i, \nu_j \rangle) &= \det \left((h^i_k)_{ik} \circ (g_{kl})_{kl} \circ (h^l_j)_{lj} \right) \\ &= \det h^i_k \cdot \det h^l_j \cdot \det g_{kl} \\ &= K^2 \cdot \det g_{kl} \end{aligned}$$

and therefore

$$A(\nu) = \int_U \sqrt{\det \langle \nu_i, \nu_j \rangle} dx = \int_U |K| \sqrt{\det g_{ij}} dx.$$

b) The image of the Gauss map covers the whole sphere S^2 . The points of the circles $(r \cos y, r \sin y, \pm a)$ are mapped to the South and North poles, respectively.

Otherwise there are exactly two points p_+ and p_- in T with the same image $q \in S^2$, one lying in the outer region T_+ with $K > 0$ and one lying in the inner region T_- with $K < 0$ (see also later). Therefore

$$\int_T K dA = \int_{T_+} K dA + \int_{T_-} K dA = A(\nu_+) - A(\nu_-) = A(S^2) - A(S^2) = 0.$$

Alternative Solution. The parametrization of the torus is given by $f: [0, 2\pi]^2 \rightarrow \mathbb{R}^3$ with

$$f(x, y) = ((r + a \cos x) \cos y, (r + a \cos x) \sin y, a \sin x),$$

where $r > a > 0$. It holds

$$\begin{aligned} f_1(x, y) &= (-a \sin x \cos y, -a \sin x \sin y, a \cos x), \\ f_2(x, y) &= (-(r + a \cos x) \sin y, (r + a \cos x) \cos y, 0), \\ f_{11}(x, y) &= (-a \cos x \cos y, -a \cos x \sin y, -a \sin x), \\ f_{12}(x, y) &= (a \sin x \sin y, -a \sin x \cos y, 0), \\ f_{22}(x, y) &= (-(r + a \cos x) \cos y, -(r + a \cos x) \sin y, 0) \end{aligned}$$

and

$$\nu(x, y) = \frac{f_1 \times f_2}{|f_1 \times f_2|} = (-\cos x \cos y, -\cos x \sin y, -\sin x).$$

From the above computations we obtain

$$\begin{aligned} (g_{ij}) &= \begin{pmatrix} a^2 & 0 \\ 0 & (r + a \cos x)^2 \end{pmatrix}, \\ (h_{ij}) &= \begin{pmatrix} a & 0 \\ 0 & (r + a \cos x) \cos x \end{pmatrix}. \end{aligned}$$

For the Gauss curvature it holds

$$K = \frac{\det(h_{ij})}{\det(g_{ij})} = \frac{a(r + a \cos x) \cos x}{a^2(r + a \cos x)^2} = \frac{\cos x}{a(r + a \cos x)}$$

and therefore

$$\int_T K \, dA = \int_0^{2\pi} \int_0^{2\pi} \frac{\cos x}{a(r + a \cos x)} \sqrt{\det(g_{ij})} \, dx \, dy = 2\pi \int_0^{2\pi} \cos x \, dx = 0.$$