

Non-Life Insurance: Mathematics and Statistics

Exercise sheet 10

Exercise 10.1 Log-Linear Gaussian Regression Model (R Exercise)

Suppose that a car insurance portfolio of an insurance company has been divided according to two tariff criteria

- vehicle type: {passenger car, delivery van, truck} = {1,2,3},
- driver age: {21-30 years, 31-40 years, 41-50 years, 51-60 years} = {1,2,3,4}.

For simplicity, we set the number of policies $v_{i,j} = 1$ for all risk classes (i, j) , $1 \leq i \leq 3$, $1 \leq j \leq 4$. Moreover, we assume that we work with a multiplicative tariff structure and that we observed the following claim amounts:

	21-30y	31-40y	41-50y	51-60y
passenger car	2'000	1'800	1'500	1'600
delivery van	2'200	1'600	1'400	1'400
truck	2'500	2'000	1'700	1'600

Table 1: Observed claim amounts in the $3 \cdot 4 = 12$ risk classes.

Calculate the tariffs using the log-linear Gaussian regression model.

- Determine the design matrix Z of the log-linear Gaussian regression model.
- Calculate the tariffs using the MLE method within the log-linear Gaussian regression model framework.
- Is there statistical evidence that the classification into different types of vehicles could be omitted?

Exercise 10.2 Method of Bailey & Simon

Consider the same setup as in Exercise 10.1. Calculate the tariffs using the method of Bailey & Simon. Comment on the results.

Exercise 10.3 Method of Bailey & Jung

Consider the same setup as in Exercise 10.1. Calculate the tariffs using the method of Bailey & Jung (i.e. the method of total marginal sums). Compare the results.

Exercise 10.4 Tweedie's Compound Poisson Model

Let $S \sim \text{CompPoi}(\lambda v, G)$, where $\lambda > 0$ is the unknown claim frequency parameter, $v > 0$ the known volume and G the distribution function of a gamma distribution with known shape parameter $\gamma > 0$ and unknown scale parameter $c > 0$. Then, S has a mixture distribution with a point mass of $\mathbb{P}[S = 0]$ in 0 and a density f_S on $(0, \infty)$.

- Calculate $\mathbb{P}[S = 0]$ and the density f_S of S on $(0, \infty)$.

(b) Show that S belongs to the exponential dispersion family with

$$\begin{aligned}
 w &= v, \\
 \phi &= \frac{\gamma + 1}{\lambda \gamma} \left(\frac{\lambda v \gamma}{c} \right)^{\frac{\gamma}{\gamma+1}}, \\
 \theta &= -(\gamma + 1) \left(\frac{\lambda v \gamma}{c} \right)^{-\frac{1}{\gamma+1}}, \\
 \Theta &= (-\infty, 0), \\
 b(\theta) &= \frac{\gamma + 1}{\gamma} \left(\frac{-\theta}{\gamma + 1} \right)^{-\gamma}, \\
 c(0, \phi, w) &= 0 \quad \text{and} \\
 c(x, \phi, w) &= \log \left(\sum_{n=1}^{\infty} \left[\frac{(\gamma + 1)^{\gamma+1}}{\gamma} \left(\frac{\phi}{w} \right)^{-\gamma-1} \right]^n \frac{1}{\Gamma(n\gamma)n!} x^{n\gamma-1} \right), \quad \text{if } x > 0.
 \end{aligned}$$

Exercise 10.5 Log-Linear Gaussian Regression Model (R Exercise)

Interpret the following R output of Exercise 10.1.

Listing 1: R output for Exercise 10.1.

```

1 Call:
2 lm(formula = observation ~ van + truck + X31_40y + X41_50y + X51_60y, data = data)
3
4 Residuals:
5      Min       1Q   Median       3Q      Max
6 -0.087095 -0.019871  0.006206  0.022773  0.064464
7
8 Coefficients:
9             Estimate Std. Error t value Pr(>|t|)
10 (Intercept)  7.68800    0.04233  181.610 1.88e-12 ***
11 van         -0.05625    0.04233  -1.329 0.232227
12 truck        0.11342    0.04233   2.679 0.036575 *
13 X31_40y     -0.21565    0.04888  -4.412 0.004511 **
14 X41_50y     -0.37511    0.04888  -7.674 0.000256 ***
15 X51_60y     -0.37381    0.04888  -7.647 0.000261 ***
16 ---
17 Signif. codes:  0 *** 0.001 ** 0.01 * 0.05 . 0.1 1
18
19 Residual standard error: 0.05987 on 6 degrees of freedom
20 Multiple R-squared:  0.941,    Adjusted R-squared:  0.8918
21 F-statistic: 19.13 on 5 and 6 DF,  p-value: 0.001261

```