

Exercise 4.1.

Prove that the Lebesgue measure is invariant under translations, reflections and rotations, i.e. under all motions of the form

$$\Phi : \mathbb{R}^n \rightarrow \mathbb{R}^n, \quad \Phi(x) = x_0 + Rx,$$

for $x_0 \in \mathbb{R}^n$ and $R \in O(n)$.

Hint: You may use the invariance of the Jordan measure, see Satz 9.3.2 in Struwe's lecture notes.

Exercise 4.2. ♣

Which of the following statements are correct?

- (a) Every countable subset of \mathbb{R} is a Borel set.
- (b) Every countable subset of \mathbb{R} is Jordan-measurable.
- (c) Every countable subset of \mathbb{R} has Lebesgue measure zero.
- (d) Every countable subset of \mathbb{R} has inner Jordan measure zero.
- (e) Every countable subset of \mathbb{R} has outer Jordan measure zero

Exercise 4.3.

(a) Let $A \subset \mathbb{R}$ be a subset with Lebesgue measure $\mathcal{L}^1(A) > 0$. Show that there exists a subset $B \subset A$ which is **not** \mathcal{L}^1 -measurable.

(b) Find an example of a countable, pairwise disjoint collection $\{E_k\}_k$ of subsets in \mathbb{R} , such that

$$\mathcal{L}^1\left(\bigcup_{k=1}^{\infty} E_k\right) < \sum_{k=1}^{\infty} \mathcal{L}^1(E_k).$$

Exercise 4.4. ★

Show that the open ball $B(x, r) := \{y \in \mathbb{R}^n \mid |y - x| < r\}$ and the closed ball $\overline{B(x, r)} := \{y \in \mathbb{R}^n \mid |y - x| \leq r\}$ in \mathbb{R}^n are Jordan measurable with Jordan measure $c_n r^n$, for some constant $c_n > 0$ depending only on n .