

Mathematical Foundations for Finance

Exercise Sheet 6

Exercise 6.1 Let $(\Omega, \mathcal{F}, \mathbb{F}, P)$ be a filtered probability space with $\mathbb{F} = (\mathcal{F}_k)_{k \in \mathbb{N}_0}$. Let $X = (X_k)_{k \in \mathbb{N}_0}$ be an adapted and integrable process.

- (a) Find the *Doob decomposition* of X . In other words, prove that there exist a martingale $M = (M_k)_{k \in \mathbb{N}_0}$ and an integrable and predictable process $A = (A_k)_{k \in \mathbb{N}_0}$ that are both null at zero, and such that

$$X = X_0 + M + A \text{ } P\text{-a.s.}$$

Hint: You may define $M_k := \sum_{j=1}^k (X_j - E[X_j | \mathcal{F}_{j-1}])$, for $k \in \mathbb{N}$.

- (b) Prove that M and A are unique up to P -a.s. equality.

Solution 6.1 To simplify notation, we omit " P -a.s." from all equalities below.

- (a) For each $k \in \mathbb{N}_0$, take

$$M_k := \sum_{j=1}^k (X_j - E[X_j | \mathcal{F}_{j-1}]).$$

It is immediate that M is adapted, integrable, and null at zero. Then, for $k \in \mathbb{N}$, we have

$$\begin{aligned} E[M_k - M_{k-1} | \mathcal{F}_{k-1}] &= E[X_k - E[X_k | \mathcal{F}_{k-1}] | \mathcal{F}_{k-1}] \\ &= E[X_k | \mathcal{F}_{k-1}] - E[X_k | \mathcal{F}_{k-1}] \\ &= 0. \end{aligned}$$

Hence, M is a martingale. Next, for each $k \in \mathbb{N}_0$, we set

$$\begin{aligned} A_k &:= X_k - X_0 - M_k = X_k - X_0 - \sum_{j=1}^k (X_j - E[X_j | \mathcal{F}_{j-1}]) \\ &= \sum_{j=1}^k (E[X_j | \mathcal{F}_{j-1}] - X_{j-1}). \end{aligned}$$

Then A is predictable with $A_0 = 0$, and of course $X = X_0 + M + A$, as required.

- (b) Suppose the processes $M^{(1)}, A^{(1)}$ and $M^{(2)}, A^{(2)}$ both satisfy the conditions of the problem. Subtracting the equalities

$$X - X_0 = M^{(1)} + A^{(1)},$$

$$X - X_0 = M^{(2)} + A^{(2)}$$

gives

$$M^{(1)} - M^{(2)} = A^{(2)} - A^{(1)}.$$

For notational convenience, we set $Y := M^{(1)} - M^{(2)} = A^{(2)} - A^{(1)}$. Since $Y = A^{(2)} - A^{(1)}$, then Y is predictable, and hence for all $k \in \mathbb{N}$,

$$Y_k = E[Y_k | \mathcal{F}_{k-1}].$$

But since the difference of two martingales is a martingale, Y is a martingale, and hence the above can be rewritten as

$$Y_k = Y_{k-1} \quad \forall k \in \mathbb{N}.$$

Since $Y_0 = 0$, this implies that $Y_k = 0$ for all $k \in \mathbb{N}_0$, and hence

$$M^{(1)} = M^{(2)} \quad \text{and} \quad A^{(1)} = A^{(2)}.$$

This completes the proof.

Exercise 6.2 Let $W = (W_t)_{t \geq 0}$ and $W' = (W'_t)_{t \geq 0}$ be two *independent* Brownian motions (BM) defined on some probability space (Ω, \mathcal{F}, P) . Show that

- $W^1 := -W$ is a BM.
- $W_t^2 := W_{T+t} - W_T$, for $t \geq 0$, is a BM for any $T \in (0, \infty)$.
- $W^3 := \alpha W + \sqrt{1 - \alpha^2} W'$ is a BM for any $\alpha \in [0, 1]$.
- Show that the independence of W and W' in (c) cannot be omitted, i.e., if W and W' are *not* independent, then W^3 need not be a BM. Give two examples.

Solution 6.2 We first recall the definition of a Brownian motion in order to know what needs to be checked. A *Brownian motion* with respect to P is a real-valued stochastic process $W = (W_t)_{t \geq 0}$ such that

(BM0) $W_0 = 0$ P -a.s.

(BM1) For any $n \in \mathbb{N}$ and any times $0 = t_0 < t_1 < \dots < t_n < \infty$, the increments $W_{t_i} - W_{t_{i-1}}$ are independent and normally distributed with variance $t_i - t_{i-1}$ under P , i.e.

$$W_{t_i} - W_{t_{i-1}} \sim \mathcal{N}(0, t_i - t_{i-1}) \quad \text{for } i = 1, \dots, n.$$

(BM2) W has P -a.s. continuous trajectories.

(a) We check (BM0), (BM1) and (BM2) separately.

(BM0) This is clear since $W_0^1 = -W_0 = 0$ P -a.s.

(BM1) Let $n \in \mathbb{N}$ and $0 = t_0 < t_1 < \dots < t_n < \infty$. Then we have, for $i = 1, \dots, n$, that

$$W_{t_i}^1 - W_{t_{i-1}}^1 = -(W_{t_i} - W_{t_{i-1}}),$$

which are independent under P . Since $X \sim \mathcal{N}(0, \sigma^2)$ if and only if $-X \sim \mathcal{N}(0, \sigma^2)$, we also conclude that $W_{t_i}^1 - W_{t_{i-1}}^1 \sim \mathcal{N}(0, t_i - t_{i-1})$.

(BM2) This is trivial, since $W^1 = -W$. The sign does not alter continuity.

(b) We check (BM0), (BM1) and (BM2) separately.

(BM0) We obviously have $W_0^2 = W_T - W_T = 0$ P -a.s.

(BM1') Let $n \in \mathbb{N}$ and $0 = t_0 < t_1 < \dots < t_n < \infty$. Then we have for $i = 1, \dots, n$ that

$$W_{t_i}^2 - W_{t_{i-1}}^2 = W_{T+t_i} - W_T - (W_{T+t_{i-1}} - W_T) = W_{T+t_i} - W_{T+t_{i-1}}.$$

Denoting $t'_i = T + t_i$, we see from the definition (BM1') that the increments of W^2 are independent under P , and since $t'_i - t'_{i-1} = t_i - t_{i-1}$, we also conclude that for all $i = 1, \dots, n$, we have

$$W_{t_i}^2 - W_{t_{i-1}}^2 \sim \mathcal{N}(0, t_i - t_{i-1}).$$

(BM2) This is again easy, since W^2 is simply W shifted in time by T minus a random variable which does not depend on t .

(c) We check (BM0), (BM1) and (BM2) separately.

(BM0) $W_0^3 = \alpha W_0 + \sqrt{1 - \alpha^2} W'_0 = 0$ P -a.s., since both W_0 and W'_0 are equal to 0 P -a.s.

(BM1') Let $n \in \mathbb{N}$ and $0 = t_0 < t_1 < \dots < t_n < \infty$. Then we have, for $i = 1, \dots, n$, that

$$W_{t_i}^3 - W_{t_{i-1}}^3 = \alpha (W_{t_i} - W_{t_{i-1}}) + \sqrt{1 - \alpha^2} (W'_{t_i} - W'_{t_{i-1}}).$$

Since W and W' are independent under P , we conclude that the right-hand side is an independent family of random variables. Since W and W' are BMs, we additionally have that

$$\begin{aligned} W_{t_i} - W_{t_{i-1}} &\sim \mathcal{N}(0, t_i - t_{i-1}), \quad i = 1, \dots, n, \\ W'_{t_i} - W'_{t_{i-1}} &\sim \mathcal{N}(0, t_i - t_{i-1}), \quad i = 1, \dots, n. \end{aligned}$$

Recall the general fact that if $X \sim \mathcal{N}(0, \sigma^2)$ and $Y \sim \mathcal{N}(0, \eta^2)$ are independent, then we have for any linear combination $s_1X + s_2Y$ that

$$s_1X + s_2Y \sim \mathcal{N}(0, s_1^2\sigma^2 + s_2^2\eta^2).$$

Using this, we conclude that

$$\alpha(W_{t_i} - W_{t_{i-1}}) + \sqrt{1 - \alpha^2}(W'_{t_i} - W'_{t_{i-1}}) \sim \mathcal{N}(0, t_i - t_{i-1})$$

since

$$\alpha^2(t_i - t_{i-1}) + (1 - \alpha^2)(t_i - t_{i-1}) = t_i - t_{i-1}.$$

(BM2) This is evident, since W^3 is a linear combination of two processes whose paths are P -a.s. continuous.

(d) Two possible choices are $W = \pm W'$. In this case, we have

$$W^3 = \left(\alpha \pm \sqrt{1 - \alpha^2}\right) W,$$

which is not a Brownian motion because $W_1^3 \sim \mathcal{N}\left(0, (\alpha \pm \sqrt{1 - \alpha^2})^2\right)$ and $(\alpha \pm \sqrt{1 - \alpha^2})^2 \neq 1$ in general.

Exercise 6.3 Let $W = (W_t)_{t \geq 0}$ be a Brownian motion defined on some probability space $(\Omega, \mathcal{F}, \mathbb{F}, P)$, where $\mathbb{F} := (\mathcal{F}_t)_{t \geq 0}$ is a filtration satisfying the usual conditions.

(a) Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be an arbitrary convex function. Show that if the stochastic process $(f(W_t))_{t \geq 0}$ is integrable, then it is a (P, \mathbb{F}) -submartingale.

Hint: We have done something similar in discrete time.

(b) Given a (P, \mathbb{F}) -martingale $(M_t)_{t \geq 0}$ and a measurable function $g: \mathbb{R}_+ \rightarrow \mathbb{R}$, show that the process

$$(M_t + g(t))_{t \geq 0}$$

is a (P, \mathbb{F}) -supermartingale if and only if g is decreasing, and a (P, \mathbb{F}) -submartingale if and only if g is increasing.

Solution 6.3

(a) First recall that W is a (P, \mathbb{F}) -martingale. Adaptedness is clear since f is continuous. Indeed, recall that any real-valued convex function is continuous on the interior of its domain. Integrability is assumed. Then by Jensen's inequality for conditional expectations, we can compute

$$E[f(W_t) | \mathcal{F}_s] \geq f(E[W_t | \mathcal{F}_s]) = f(W_s) \text{ } P\text{-a.s.}$$

for all $t \geq s$, and thus conclude that $f(W)$ is a (P, \mathbb{F}) -submartingale.

- (b) For any measurable function $g : \mathbb{R}_+ \rightarrow \mathbb{R}$, we have that, for each $t \geq 0$, $M_t + g(t)$ is \mathcal{F}_t -measurable and

$$E[|M_t + g(t)|] \leq E[|M_t|] + E[|g(t)|] = E[|M_t|] + |g(t)| < \infty.$$

Hence $(M_t + g(t))_{t \geq 0}$ is adapted and integrable. We can then compute

$$E[M_t + g(t) | \mathcal{F}_s] = E[M_t | \mathcal{F}_s] + g(t) = M_s + g(s) + g(t) - g(s) \text{ } P\text{-a.s.}$$

for all $t \geq s$. As a result, $(M_t + g(t))_{t \geq 0}$ has the (P, \mathbb{F}) -supermartingale property, i.e.

$$E[M_t + g(t) | \mathcal{F}_s] \leq M_s + g(s) \text{ } P\text{-a.s.}$$

for all $t > s$, if and only if g is decreasing. Analogously, $(M_t + g(t))_{t \geq 0}$ has the (P, \mathbb{F}) -submartingale property, i.e.

$$E[M_t + g(t) | \mathcal{F}_s] \geq M_s + g(s) \text{ } P\text{-a.s.}$$

for all $t > s$, if and only if g is increasing.