

Mathematical Foundations for Finance

Exercise Sheet 9

Please hand in your solutions by 12:00 on Wednesday, November 29 via the course homepage.

Exercise 9.1 On a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, P)$, consider an adapted process $X = (X_t)_{t \geq 0}$ null at 0. Assume that X is integrable and has independent and stationary increments, i.e. $X_t - X_s$ is independent of \mathcal{F}_s and has the same distribution as X_{t-s} for all $t > s \geq 0$.

- (a) Under which conditions on $(E[X_t])_{t \geq 0}$ is X a martingale? And a supermartingale? A submartingale?
- (b) From this point onward, let us assume that X is a square-integrable martingale. Prove that

$$E[X_t^2] + E[X_s^2] = E[X_{t+s}^2] \text{ for any } t, s \geq 0,$$

and deduce that $(E[X_t^2])_{t \geq 0}$ is an increasing process.

Hint: You may use Exercise 8.1(a).

- (c) Deduce from (b) that $E[X_t^2] = tE[X_1^2]$ for all $t \geq 0$.
Hint: Prove the result first for $t = 1/n$ for all $n \in \mathbb{N}$. Then, deduce that it holds true for all $t \in \mathbb{Q}_+$ and use monotonicity to conclude.
- (d) Prove that $\langle X \rangle_t = tE[X_1^2]$ for all $t \geq 0$.
Hint: You may use your result from (c).

Exercise 9.2 Consider a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, P)$, where the filtration \mathbb{F} satisfies the usual conditions.

- (a) Let X be an adapted process and τ a stopping time. Show that if X^τ is a martingale, then so is X^σ for any stopping time σ with $\sigma \leq \tau$ P -a.s.
Hint: You may use the result that a stopped martingale is again a martingale.
- (b) Let M and N be two local martingales. Show that the linear combination $\alpha M + \beta N$ for any $\alpha, \beta \in \mathbb{R}$ is a local martingale.
Hint: You may use your result in (a).
- (c) We say that two Brownian motions W^1 and W^2 on $(\Omega, \mathcal{F}, \mathbb{F}, P)$ are *correlated with instantaneous correlation* $\rho \in [-1, 1]$ if, for $s \leq t$, the increments $W_t^1 - W_s^1$

and $W_t^2 - W_s^2$ are independent of \mathcal{F}_s and jointly normally distributed with $\mathcal{N}(\mu, \Sigma)$, where

$$\mu = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \text{ and } \Sigma = \begin{pmatrix} t-s & \rho(t-s) \\ \rho(t-s) & t-s \end{pmatrix}.$$

Show that $[W^1, W^2]_t = \rho t$ P -a.s.

Hint: You may find $\lambda \in \mathbb{R}$ such that $B^\lambda := \lambda(W^1 + W^2)$ is a Brownian motion. Then, compute $[B^\lambda]$ in terms of W^1 and W^2 , using the properties of $[\cdot, \cdot]$.

Exercise 9.3 Let $(\Omega, \mathcal{F}, \mathbb{F}, P)$ be a filtered probability space, where the filtration \mathbb{F} satisfies the usual conditions. Consider $W = (W_t)_{t \geq 0}$ and $B = (B_t)_{t \geq 0}$ two independent Brownian motions.

- (a) Show that $BW = (B_t W_t)_{t \geq 0}$ is a martingale.
- (b) Compute the mean of $\int_0^t W_s ds$ and prove that its variance is $t^3/3$, for $t > 0$.
Hint: You may use Fubini's theorem, and rewrite $(\int_0^t W_s ds)^2 = \int_0^t \int_0^t W_s W_u ds du$.