

Solutions to problem set 4

1. Consider the commutative diagram

$$\begin{array}{ccc} H_0(X) & \xrightarrow{(\epsilon_X)_*} & H_0(P) \\ \downarrow f_* & & \downarrow id \\ H_0(Y) & \xrightarrow{(\epsilon_Y)_*} & H_0(P). \end{array}$$

Here P denotes the one-point space and $(\epsilon_X)_*$ is the homomorphism induced by the unique map $\epsilon_X: X \rightarrow P$. Similarly for ϵ_Y . It follows from a simple diagram chase that f_* induces a homomorphism

$$f_*: \tilde{H}_0(X) = \ker((\epsilon_X)_*) \rightarrow \ker((\epsilon_Y)_*) = \tilde{H}_0(Y)$$

indeed if $\alpha \in \ker((\epsilon_X)_*)$, then

$$(\epsilon_Y)_*(f_*\alpha) = f_*((\epsilon_X)_*\alpha) = f_*(0) = 0.$$

2. Let $1 \leq i \neq j \leq n$. Consider the following rotations $R_t^{i,j}$ in \mathbb{R}^{n+1} : In the x_i - x_j -plane, $R_t^{i,j}$ is represented by the matrix

$$R_t^{i,j} = \begin{pmatrix} \cos(\frac{\pi}{2}t) & \sin(\frac{\pi}{2}t) \\ -\sin(\frac{\pi}{2}t) & \cos(\frac{\pi}{2}t) \end{pmatrix}.$$

$R_t^{i,j}$ fixes the other coordinates x_k , $k \neq i, j$. Then $R_t^{i,j}$ restrict to homeomorphisms on S^n and $\tau_i = (R_1^{i,j})^{-1} \circ \tau_j \circ R_1^{i,j}$. Thus $\left\{ (R_t^{i,j})^{-1} \circ \tau_j \circ R_t^{i,j} \right\}_{t \in [0,1]}$ is a homotopy from τ_j to τ_i .

3. We show that \hat{f} is continuous: Let $V \subset \hat{Y}$ be an open subset. If $V \subset Y$ is open, then $\hat{f}^{-1}(V) = f^{-1}(V) \subset X$ is open in X by continuity of f . Thus $\hat{f}^{-1}(V)$ is open in \hat{X} . If $\infty \in V$, then $\infty \in \hat{f}^{-1}(V)$ and $\hat{X} \setminus \hat{f}^{-1}(V) = f^{-1}(\hat{Y} \setminus V)$. Note that $\hat{Y} \setminus V \subset Y$ is compact. Since f is a homeomorphism, f is proper and so $f^{-1}(\hat{Y} \setminus V)$ is compact. It follows that $\hat{f}^{-1}(V)$ is open in \hat{X} . The same argument applied to f^{-1} implies that $\widehat{f^{-1}}$ is continuous. Thus \hat{f} is a homeomorphism with inverse $\widehat{f^{-1}}$.

If we drop the assumption that f is a homeomorphism, then f doesn't necessarily extend to a continuous map \hat{f} on the compactifications. Here is a counterexample: Consider the inclusion i of the 1-disk (i.e. the open unit interval) $B_1(0)$ into the 2-disk $B_2(0)$ as an equator. Then there is no continuous extension of i to the compactifications: indeed the subspace

$$V = \left(\widehat{B_2(0)} \setminus \overline{B_1(0)} \right) \cup \{\infty\}$$

is open in $\widehat{B_2(0)}$ as $\overline{B_1(0)}$ is compact, while its inverse image $\hat{i}^{-1}(V) = \{\infty\} \in \widehat{B_1(0)}$ is not open, as $\{\infty\} = \overline{B_1(0)} \setminus B_1(0)$, but $B_1(0)$ is not compact.

One can show that a map extends continuously to one-point compactifications if and only if it is a proper map.

4. View S^n as the standard sphere in \mathbb{R}^{n+1} with coordinates (x_0, x_1, \dots, x_n) . Define the stereographic projection $\pi: S^n \setminus \{(1, 0, \dots, 0)\} \rightarrow \mathbb{R}^n$ as follows:

$$\pi(x_0, x_1, \dots, x_n) = \left(\frac{x_1}{1-x_0}, \dots, \frac{x_n}{1-x_0} \right).$$

$\pi(x)$ is the intersection of the unique line through x and $(1, 0, \dots, 0)$ with the hyperplane $\{x_0 = 0\}$. π is a homeomorphism with inverse

$$\pi^{-1}(y_1, \dots, y_n) = \left(\frac{\|y\|^2 - 1}{\|y\|^2 + 1}, \frac{2y_1}{\|y\|^2 + 1}, \dots, \frac{2y_n}{\|y\|^2 + 1} \right).$$

It follows from Exercise 3 that π extends to a homeomorphism $\hat{\pi}: S^n \rightarrow \mathbb{R}^n \cup \{\infty\}$.

5. View S^{2k-1} as the unit sphere inside \mathbb{C}^k , with respect to the standard Euclidean metric on \mathbb{C}^k . For every point $z \in S^{2k-1}$, viewed as a k -tuple of complex numbers, consider the curve $\gamma_z: (-\epsilon, \epsilon) \rightarrow S^{2k-1}$ given by $\gamma_z(t) = e^{it}z$. Consider the vector field X on S^{2k-1} given by

$$X(z) = \dot{\gamma}_z(0).$$

This is a smooth nowhere vanishing vector field. In real coordinates it is given by

$$X(x_1, y_1, \dots, x_k, y_k) = (-y_1, x_1, \dots, -y_k, x_k).$$

6. (a) Since $f(x) \neq x, \forall x \in S^n$, the line segment $(1-t)f(x) - tx, t \in [0, 1]$, does not pass through 0. Therefore, if f has no fixed points,

$$f_t(x) := \frac{(1-t)f(x) - tx}{|(1-t)f(x) - tx|}$$

is a well defined homotopy from f to the antipodal map $-id$ which has degree $\deg(-id) = (-1)^{n+1}$. Thus $\deg f = (-1)^{n+1}$.

- (b) Since $\deg f = 0 \neq (-1)^{n+1}$ it must have a fixed point $x \in S^n$ by exercise 6.(a), i.e. $f(x) = x$. Similarly, since $g := (-id) \circ f$ has degree $\deg g = \deg(-id) \cdot \deg f = 0$, there is a fixed point $y \in S^n$ of g , i.e. $g(y) = -f(y) = y$. This means that $f(y) = -y$.
7. See Example 2.32 on page 137 in Hatcher's book, or the notes from the exercise class of the 15.11.2024.
8. (a) Recall from Exercise 4 in Problem set 3 that we have the following commutative diagram

$$\begin{array}{ccc} \tilde{H}_{n+1}(S^{n+1}) & \xrightarrow[\cong]{\partial_*} & \tilde{H}_n(S^n) \\ \downarrow (Sf)_* & & \downarrow f_* \\ \tilde{H}_{n+1}(S^{n+1}) & \xrightarrow[\cong]{\partial_*} & \tilde{H}_n(S^n) \end{array}$$

(i.e. ∂_* is natural). Therefore, if f_* is multiplication by $d = \deg f$, then $(Sf)_*$ is also multiplication by d and hence $\deg f = \deg Sf$.

- (b) Given $k \in \mathbb{Z}$ the map $S^1 \rightarrow S^1: z \mapsto z^k$ has degree k . Now assume that we have constructed a map $f: S^n \rightarrow S^n$ of degree k , then (by exercise 8.(a)), the map $Sf: S^{n+1} \rightarrow S^{n+1}$ has degree k as well. So the claim follows by induction.

9. First, let $n = 1$ and denote $I := [0, 1]$. Let $g : I \rightarrow \mathbb{R}$ be a continuous map such that $g(0) = g(1) = 0$ and $g(1/2) = 2\pi$. The map g induces a well defined continuous surjection $f : I/\partial I = S^1 \rightarrow S^1 : t \mapsto e^{ig(t)}$. By the path lifting property the map g is the unique lift of f to the universal cover \mathbb{R} of S^1 starting at the point $0 \in \mathbb{R}$. So $f \in p_{\#} \underbrace{\pi_1(\mathbb{R}, 0)}_{=0} \subset \pi_1(S^1, 1)$ is homotopic to a constant map (which is clearly not surjective and therefore has degree 0) and hence $\deg f = 0$. Here, $p : \mathbb{R} \rightarrow S^1$ is the universal cover.

Using exercise 8.(a) we obtain, by repeatedly suspending the map f , a surjective map $S^n \rightarrow S^n$ of degree 0.

For an alternative, more explicit, solution see example 2.31 in Hatcher's book.

10. Let the group action be given by the homomorphism $\rho : G \rightarrow \text{Homeo}(S^n)$. The degree of a homeomorphism is always ± 1 . Therefore the group action determines a degree function $d : G \rightarrow \{\pm 1\}$ given by $d(g) := \deg \rho(g)$. Furthermore d is a homomorphism:

$$d(hg) = \deg \rho(hg) = \deg(\rho(h) \circ \rho(g)) = \deg \rho(h) \cdot \deg \rho(g) = d(h) \cdot d(g).$$

If $g \in G$ is a non trivial element, then $\rho(g)$ has no fixed points as the action is free and hence (by exercise 6.(a)) we have $d(g) = (-1)^{n+1}$. So, if n is even, the kernel of d is trivial which implies that G is isomorphic to a subgroup of $\{\pm 1\} \cong \mathbb{Z}_2$.

11. For $n = 2$ we have that $SO(2)$ is homeomorphic to the circle S^1 which is path connected. Proceeding by induction we assume that $SO(n-1)$ is path connected. Given any $A \in SO(n)$ it is enough to show that there is a path in $SO(n)$ connecting A to the identity matrix I_n . This means that we need to find a continuous path taking the standard basis e_1, \dots, e_n to their image Ae_1, \dots, Ae_n . Let $\Lambda \subset \mathbb{R}^n$ be a plane containing both e_1 and Ae_1 . By the path connectedness of $SO(2)$, we can continuously move e_1 to Ae_1 by a rotation R of the plane Λ .

It remains to continuously move Re_2, \dots, Re_n to Ae_2, \dots, Ae_n while keeping Ae_1 fixed. Notice that $Ae_1 = Re_1 \perp Re_i$ and $Ae_1 \perp Ae_i$ for each $2 \leq i \leq n$ since both R and A preserve angles. Hence the required motion can take place in the hyperplane \mathbb{R}^{n-1} of vectors orthogonal to Ae_1 , where it exists by the assumption that $SO(n-1)$ is path connected.

Concatenating the two motions gives a path in $SO(n)$ from I_n to A and thus $SO(n)$ is path connected.

For the other groups, take a look at

<https://www.jnu.ac.in/Faculty/vedgupta/matrix-gps-gupta-mishra.pdf>