

Problem set 4

1. Let H be any homology theory. Let X and Y be two non-empty spaces and $f: X \rightarrow Y$ a map. Show that the homomorphism $f_*: H_0(X) \rightarrow H_0(Y)$ induces a homomorphism $\tilde{f}_*: \tilde{H}_0(X) \rightarrow \tilde{H}_0(Y)$.

2. For $1 \leq i \leq n$ consider the maps

$$\begin{aligned} \tau_i: S^n &\longrightarrow S^n \\ (x_1, \dots, x_{n+1}) &\longmapsto (x_1, \dots, -x_i, \dots, x_{n+1}). \end{aligned}$$

Show that any two of these maps are homotopic.

3. For a space X , denote by \hat{X} its 1-point compactification. Show that if $f: X \rightarrow Y$ is a homeomorphism then f induces a homeomorphism $\hat{f}: \hat{X} \rightarrow \hat{Y}$ with $\hat{f}(\infty) = \infty$ and $\hat{f}|_X = f$. What happens if we drop the assumption that f is a homeomorphism?
4. Show that the stereographic projection $\hat{\pi}: S^n \rightarrow \hat{\mathbb{R}}^n = \mathbb{R}^n \cup \{\infty\}$ is a homeomorphism. Derive a formula for $\hat{\pi}$.
5. Construct a nowhere vanishing vector field on the odd-dimensional sphere S^{2k-1} .
Hint: View S^{2k-1} as the unit sphere inside \mathbb{C}^k , with respect to the standard Euclidean metric on \mathbb{C}^k . For every point $z \in S^{2k-1}$, viewed as a k -tuple of complex numbers, consider the curve $\gamma_z: (-\epsilon, \epsilon) \rightarrow S^{2k-1}$ given by $\gamma_z(t) = e^{it}z$.
6. (a) If $f: S^n \rightarrow S^n$ has no fixed points, then $\deg f = (-1)^{n+1}$.
(b) Let $f: S^n \rightarrow S^n$ be a map of degree 0. Show that there exist points $x, y \in S^n$ such that $f(x) = x$ and $f(y) = -y$.
7. We view S^1 as the unit circle in \mathbb{C} . Show that for each $k \in \mathbb{Z}$ the map $f: S^1 \rightarrow S^1$, given by $f(z) = z^k$, has degree k .
8. (a) We denote by SX the (unreduced) suspension of a space X we introduced in Exercise 4 in Problem set 3. If $X = S^n$, then $SX \approx S^{n+1}$. Let $Sf: S^{n+1} \rightarrow S^{n+1}$ be the suspension of a map $f: S^n \rightarrow S^n$. Show that $\deg Sf = \deg f$.
(b) Show that for each $n \geq 1$ and each $k \in \mathbb{Z}$ there is a map $f: S^n \rightarrow S^n$ of degree k .

9. Construct a surjective map $S^n \rightarrow S^n$ of degree 0 for each $n \geq 1$.
Hint: Do it first for $n = 1$ and then use exercise 8(a).
10. Let G be a group acting freely on S^n . Prove that if n is even, then G is either trivial or isomorphic to $\mathbb{Z}_2 := \mathbb{Z}/2\mathbb{Z}$.
11. Consider $SO(n)$ as a subset of \mathbb{R}^{n^2} and endow $SO(n)$ with the induced topology. Show that $SO(n)$ is path connected. Similarly, show that the Lie groups $GL(n, \mathbb{C})$, $GL^+(n, \mathbb{R})$, $U(n)$, $SU(n)$ are path connected.