

## Solutions to problem set 6

- Denote by  $D_1, D_2, D_3 \subset X$  the images of the three discs that get glued together. With  $A := D_1 \cup D_2$  and  $B := D_2 \cup D_3$ , we have  $X = A \cup B$ . We will apply Mayer-Vietoris to  $(X, A, B)$ . (The fact that the interiors of  $A$  and  $B$  do *not* cover  $X$  does not cause a problem here, because  $A$  and  $B$  have open neighbourhoods  $A'$  and  $B'$  which deformation retract onto them, obtained by adding a small “collar” neighbourhood to each, and such that  $A' \cap B'$  deformation retracts onto  $A \cap B$ . Convince yourself of that!).

$A$  and  $B$  are homeomorphic to  $S^n$ , and  $A \cap B = D_2$  is a copy of  $D^n$ . Hence  $H_*(A \cap B)$  vanishes except in degree 0, and  $H_*(A), H_*(B)$  vanish except in degrees 0 and  $n$ . This implies that the third arrow in the following portion of the Mayer-Vietoris sequence is an isomorphism for all  $k \geq 2$ :

$$0 \rightarrow H_k(A \cap B) \rightarrow H_k(A) \oplus H_k(B) \rightarrow H_k(X) \rightarrow H_{k-1}(A \cap B) \rightarrow \dots$$

i.e.,  $H_k(X) \cong H_k(A) \oplus H_k(B)$  for all  $k \geq 2$ . For  $k = 1$ , we obtain

$$0 \rightarrow H_1(A) \oplus H_1(B) \rightarrow H_1(X) \rightarrow H_0(A \cap B) \rightarrow H_0(A) \oplus H_0(B) \rightarrow H_0(X) \rightarrow 0$$

Note that  $H_0(A \cap B) \rightarrow H_0(A) \oplus H_0(B)$  is injective as  $A \cap B = D_2$  is path-connected, and hence  $H_1(X) \rightarrow H_0(A \cap B)$  is zero by exactness. Hence we obtain an isomorphism  $H_1(X) \cong H_1(A) \oplus H_1(B)$  also in this case. Moreover, it is clear that  $H_0(X) \cong \mathbb{Z}$  as  $X$  is path-connected. Putting all together, we obtain

$$H_k(X) = \begin{cases} \mathbb{Z}, & k = 0, \\ \mathbb{Z}^2, & k = n, \\ 0, & \text{otherwise.} \end{cases}$$

- We view  $\mathbb{R}P^2$  as  $D^2 / \sim$ , the quotient of  $D^2$  obtained by identifying antipodal points on  $S^1 = \partial D^2$ . Let  $A \subset \mathbb{R}P^2$  be the image of the interior of  $D^2$  under the projection  $D^2 \rightarrow \mathbb{R}P^2$ , and let  $B \subset \mathbb{R}P^2$  be the image of a collar neighbourhood of  $\partial D^2$ , homeomorphic to  $[0, \varepsilon) \times \partial D^2$ . The subsets form an open cover of  $\mathbb{R}P^2$ , and  $A \cap B \simeq S^1$ . The Mayer-Vietoris sequence for  $(\mathbb{R}P^2, A, B)$  is

$$\begin{aligned} 0 \rightarrow H_2(\mathbb{R}P^2) \rightarrow H_1(A \cap B) \rightarrow H_1(A) \oplus H_1(B) \rightarrow H_1(\mathbb{R}P^2) \\ \rightarrow H_0(A \cap B) \rightarrow H_0(A) \oplus H_0(B) \rightarrow H_0(\mathbb{R}P^2) \rightarrow 0, \end{aligned}$$

Note that  $H_0(A \cap B) \rightarrow H_0(A) \oplus H_0(B)$  is injective since  $A \cap B \simeq S^1$  is path-connected; moreover  $H_1(A \cap B) = \mathbb{Z} = H_1(B)$ , and  $H_1(A) = 0$ . Hence we obtain an exact sequence

$$0 \rightarrow H_2(\mathbb{R}P^2) \rightarrow \mathbb{Z} \rightarrow \mathbb{Z} \rightarrow H_1(\mathbb{R}P^2) \rightarrow 0.$$

The middle map  $\mathbb{Z} \rightarrow \mathbb{Z}$  is multiplication by 2 because the canonical map  $D^2 \rightarrow \mathbb{R}P^2$  restricts to a degree 2 map on  $\partial D^2$ . It follows that  $H_2(\mathbb{R}P^2) = 0$  and  $H_1(\mathbb{R}P^2) = \mathbb{Z}_2$ ; moreover  $H_0(\mathbb{R}P^2) = \mathbb{Z}$  as  $\mathbb{R}P^2$  is path-connected.

3. Let  $A \subset K$  be the image of the interior of  $I^2$  under the projection  $I^2 \rightarrow K$ , and let  $B \subset K$  be the image of a neighbourhood of  $\partial I^2$  homeomorphic to  $[0, \varepsilon) \times \partial I^2$ . Note that  $A$  is homeomorphic to  $D^2$ , while  $B$  deformation retracts onto the image of  $\partial I^2$  in  $K$ , which is homeomorphic to  $S^1 \vee S^1$ . The intersection  $A \cap B$  is homotopy equivalent to  $S^1$ . By the same arguments as in the previous problem, the Mayer-Vietoris sequence for  $(K, A, B)$  yields an exact sequence of the form

$$0 \rightarrow H_2(K) \rightarrow H_1(A \cap B) \rightarrow H_1(B) \rightarrow H_1(K) \rightarrow 0$$

For suitable identifications  $H_1(A \cap B) \cong \mathbb{Z}$  and  $H_1(B) \cong \mathbb{Z}^2$ , the middle map is  $\mathbb{Z} \xrightarrow{(0,2)} \mathbb{Z}^2$ . It follows that  $H_2(K) = 0$ ,  $H_1(K) = \mathbb{Z}_2 \oplus \mathbb{Z}$ ; moreover,  $H_0(K) = \mathbb{Z}$  as  $K$  is path-connected.

4. Denote by  $P$  our polygon and by  $p : P \rightarrow \Sigma_g$  the canonical projection. Let  $A$  be the image of the interior of  $P$  under  $p$ , and let  $B$  be the image of a neighbourhood of  $\partial P$  which is homeomorphic to  $[0, \varepsilon) \times S^1$ . From the Mayer-Vietoris sequence for  $(\Sigma_g, A, B)$  we obtain an exact sequence

$$0 \rightarrow H_2(\Sigma_g) \rightarrow H_1(A \cap B) \rightarrow H_1(B) \rightarrow H_1(\Sigma_g) \rightarrow 0$$

by the same argument as in the previous problems. The image of  $\partial P$  under  $p$ , considered as a loop in  $\Sigma_g$ , represents the class  $[a_1][b_1][a_1^{-1}][b_1^{-1}] \dots [a_g][b_g][a_g^{-1}][b_g^{-1}]$  in  $\pi_1(B)$ , which lies in the commutator subgroup of  $\pi_1(B)$ ; hence its image in  $H_1(B)$  vanishes. The map  $H_1(A \cap B) \rightarrow H_1(B)$  is therefore zero, and we obtain  $H_2(\Sigma_g) \cong H_1(A \cap B) \cong \mathbb{Z}$  and  $H_1(\Sigma_g) \cong H_1(B) \cong \mathbb{Z}^{2g}$  by exactness and because  $A \cap B \simeq S^1$  and  $B \simeq S^1 \vee \dots \vee S^1$  ( $2g$  times).  $H_0(\Sigma_g) \cong \mathbb{Z}$  is clear because  $\Sigma_g$  is path-connected.

5. We will show by induction that  $H_0(\Sigma_g) \cong \mathbb{Z}$ ,  $H_1(\Sigma_g) \cong \mathbb{Z}^{2g}$ , and  $H_2(\Sigma_g) \cong \mathbb{Z}$ . For  $\Sigma_1 = T^2$ , this can be shown using e.g. cellular homology. To show it holds for  $\Sigma_{g+1}$  (assuming it's already shown for  $\Sigma_k$ ,  $k = 1, \dots, g$ ), consider the cover of  $\Sigma_{g+1}$  by open subsets  $A$  and  $B$ , where  $A$  and  $B$  are homeomorphic to  $\Sigma_g^* := \Sigma_g \setminus \{\text{pt}\}$  resp.  $\Sigma_1^* := \Sigma_1 \setminus \{\text{pt}\}$ ; the existence of such a cover is indicated by the definition of  $\Sigma_{g+1}$  as the connected sum of  $\Sigma_g$  and  $\Sigma_1$ .

The punctured surface  $\Sigma_g^*$  is homotopy equivalent to  $S^1 \vee \dots \vee S^1$ , a wedge of  $2g$  circles (one can see this using the polygon description of  $\Sigma_g$  given in problem 5.5), and hence  $H_2(\Sigma_g^*) \cong 0$ ,  $H_1(\Sigma_g^*) \cong \mathbb{Z}^{2g}$  as one can see using e.g. cellular homology. Since moreover the intersection  $A \cap B$  is homotopy equivalent to  $S^1$ , the Mayer-Vietoris sequence for  $(\Sigma_{g+1}, A, B)$  is

$$0 \rightarrow H_2(\Sigma_{g+1}) \rightarrow H_1(A \cap B) \rightarrow H_1(A) \oplus H_1(B) \rightarrow H_1(\Sigma_{g+1}) \rightarrow H_0(A \cap B) \rightarrow \dots \quad (1)$$

The homomorphism  $H_1(A \cap B) \rightarrow H_1(A) \oplus H_1(B)$  is zero because  $A \cap B$  deformation retracts onto a loop which lies in the commutator subgroup of both  $\pi_1(A)$  and  $\pi_1(B)$  (using the polygon description, see the solution of problem 5.5); the homomorphism  $H_1(\Sigma_{g+1}) \rightarrow H_0(A \cap B)$  is zero because  $A \cap B$  is path-connected, and hence the inclusions of  $S^1$  into  $A$  and  $B$  induce injective maps on  $H_0$ . We therefore obtain isomorphisms  $H_2(\Sigma_{g+1}) \cong H_1(S^1) \cong \mathbb{Z}$  and  $H_1(\Sigma_{g+1}) \cong H_1(\Sigma_g^*) \oplus H_1(\Sigma_1^*) \cong \mathbb{Z}^{2g} \oplus \mathbb{Z}^2 = \mathbb{Z}^{2g+2}$ , as required.  $H_0(\Sigma_{g+1}) \cong \mathbb{Z}$  is clear as  $\Sigma_{g+1}$  is path-connected.


*(Remark.* Using the polygon description of  $\Sigma_g$  is a bit of a short-cut, with which one could dispense as follows. First one computes inductively that  $H_2(\Sigma_g^*) = 0$  and  $H_1(\Sigma_g^*) = \mathbb{Z}^{2g}$ , viewing  $\Sigma_{g+1}$  as the connected sum of  $\Sigma_g$  and  $\Sigma_1^*$  and applying Mayer-Vietoris to  $(\Sigma_{g+1}^*, \Sigma_g^*, \Sigma_1^{**})$ , where  $\Sigma_1^{**}$  denotes a twice-punctured torus. To get the induction started, one computes that  $H_2(\Sigma_1^*) = 0 = H_2(\Sigma_1^{**})$ ,  $H_1(\Sigma_1^*) = \mathbb{Z}^2$  and  $H_1(\Sigma_1^{**}) = \mathbb{Z}^3$  using cellular homology. The Mayer-Vietoris sequence for  $(\Sigma_{g+1}^*, \Sigma_g^*, \Sigma_1^{**})$  yields an exact sequence

$$0 \rightarrow H_2(\Sigma_{g+1}^*) \rightarrow H_1(S^1) \rightarrow H_1(\Sigma_g^*) \oplus H_1(\Sigma_1^{**}) \rightarrow H_1(\Sigma_{g+1}^*) \rightarrow 0.$$

The map  $H_1(S^1) \rightarrow H_1(\Sigma_g^*) \oplus H_1(\Sigma_1^{**})$  is injective because the intersection  $\Sigma_g^* \cap \Sigma_1^{**} \simeq S^1$  generates a  $\mathbb{Z}$ -summand in  $H_1(\Sigma_1^{**})$  (check this using cellular homology), and hence the second component of the map is non-zero. This implies that  $H_2(\Sigma_{g+1}^*) = 0$  and  $H_1(\Sigma_{g+1}^*) \cong \mathbb{Z}^{2g+2}$ , using the inductive hypothesis. Having computed  $H_*(\Sigma_g^*)$ , one proceeds with the inductive computation of  $H_*(\Sigma_{g+1}^*)$  using the Mayer-Vietoris sequence (??); the vanishing of  $H_1(A \cap B) \rightarrow H_1(A) \oplus H_1(B)$  can be checked by constructing inductively 2-chains in  $A$  and  $B$  whose boundary is a circle homotopy equivalent to  $A \cap B$ .)

6. For  $S^0 = \{1, -1\}$ , set  $c_0 := 1 - (-1) \in \Delta_0(S^0)$ . Then  $[c_0] = [1] - [-1]$  generates  $\tilde{H}_0(S^0)$ . (Recall that  $\tilde{H}_0(S^0)$  is the kernel of the canonical map  $H_0(S^0) \rightarrow H_0(\text{pt})$ , which maps both generators  $[-1]$  and  $[1]$  of  $H_0(S^0)$  to  $[\text{pt}]$ .)

As for  $S^1$ , consider 1-simplices  $c_1^0, c_1^1 : \Delta_1 \rightarrow S^1$  as indicated in the figure below, and let  $c_1 := c_1^0 + c_1^1 \in \Delta_1(S^1)$ , which is a cycle. Consider now the cover of  $S^1$  by  $A = \text{im } c_1^0$  and  $B = \text{im } c_1^1$  and the reduced Mayer-Vietoris sequence for  $(S^1, A, B)$  (which exists despite the fact that the interiors of  $A$  and  $B$  do not cover  $S^1$ , by the same reason as in problem 5.2). Then  $A \cap B = S^0$ , and the boundary morphism  $\partial_* : \tilde{H}_1(S^1) \rightarrow \tilde{H}_0(A \cap B) = \tilde{H}_0(S^0)$  takes  $[c_1]$  to  $[\partial c_1^0] = [1 - (-1)] = [c_0]$ , our generator of  $\tilde{H}_0(S^0)$ .



To find a generator of  $\tilde{H}_2(S^2)$ , view  $S^2$  as the union of two copies  $D_-$  and  $D_+$  of  $D^2$ , identified along their boundaries. Consider simplices  $c_2^0, c_2^1, c_2^2, c_2^3 : \Delta_2 \rightarrow S^2$  as indicated in the figure; note that  $c_2 = c_2^0 + c_2^1 + c_2^2 + c_2^3 \in \Delta_2(S^2)$  is a cycle. Consider now the Mayer-Vietoris sequence for  $(S^2, D_-, D_+)$ . We have  $D_- \cap D_+ = S^1$ , and the boundary operator  $\partial_* : H_2(S^2) \rightarrow H_1(D_- \cap D_+) = H_1(S^1)$  maps  $[c_2]$  to  $[\partial c_2^0 + \partial c_2^1] \in H_1(D_- \cap D_+)$ , which is precisely  $[c_1^0 + c_1^1] = [c_1]$ .

7. Set  $A = X \cup V$  and  $B = Y \cup U$ , thinking of these as subsets of  $X \vee Y$ . Then  $A$  deformation retracts onto  $X$ ,  $B$  deformation retracts onto  $Y$ , and  $A \cap B$  deformation retracts onto  $\{*\}$ , the point at which the two spaces are joint. The reduced Mayer-Vietoris sequence for  $(X \vee Y, A, B)$  takes the form

$$\cdots \rightarrow \tilde{H}_k(\{*\}) \rightarrow \tilde{H}_k(X) \oplus \tilde{H}_k(Y) \rightarrow \tilde{H}_k(X \vee Y) \rightarrow \tilde{H}_{k-1}(\{*\}) \rightarrow \cdots,$$

which yields  $\tilde{H}_k(X \vee Y) \cong \tilde{H}_k(X) \oplus \tilde{H}_k(Y)$  for all  $k$ , because  $\tilde{H}_k(\{*\}) = 0$  for all  $k$ .

8. Let  $A, B \subset SX$  be the images of  $[0, 1) \times X$  resp.  $(0, 1] \times X$  under the canonical map  $X \times I \rightarrow SX$ .  $A$  and  $B$  are both contractible and hence have vanishing reduced homology, while  $A \cap B \simeq X$ . The reduced Mayer-Vietoris sequence

$$\cdots \rightarrow \tilde{H}_k(A) \oplus \tilde{H}_k(B) \rightarrow \tilde{H}_k(SX) \rightarrow \tilde{H}_{k-1}(X) \rightarrow \tilde{H}_{k-1}(A) \oplus \tilde{H}_{k-1}(B) \rightarrow \cdots$$

hence yields  $\tilde{H}_k(SX) \cong \tilde{H}_{k-1}(X)$  for all  $k$ .