

# PROBABILITY AND STATISTICS

## Exercise sheet 4

**MC 4.1. [Even-indexed Poisson(1)]** Let  $X$  be a random variable taking values in  $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$  with

$$\mathbb{P}[X = k] = \begin{cases} c/k!, & k \in \mathbb{N}_0, k \text{ even,} \\ 0, & \text{otherwise,} \end{cases}$$

where, for simplicity, we consider 0 as an even number. For which values of  $c \in \mathbb{R}$  does this define a distribution? (Exactly one answer is correct.)

**Hint:** You may use the following identity:

$$\sum_{n=0}^{\infty} \frac{1}{(2n)!} = \sum_{n=0}^{\infty} \frac{1}{2} \left( \frac{1}{n!} + \frac{(-1)^n}{n!} \right) = \frac{1}{2} \left( \sum_{n=0}^{\infty} \frac{1}{n!} + \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} \right).$$

- (a) Never.
- (b) Only for  $c = e^{-1}$ .
- (c) For all  $c \geq 0$ .
- (d) Only for  $c = 2/(e + e^{-1})$ .

**MC 4.2.** Let  $X$  be a random variable taking values in  $\{1, \dots, 10\}$  with  $\mathbb{P}[X = k] = k - c$  for  $k \in \{1, \dots, 10\}$ . For which values of  $c \in \mathbb{R}$  does this define a distribution? (Exactly one answer is correct.)

- (a) If  $c = 5.4$ .
- (b) Never.
- (c) If  $c = 1$ .
- (d) If  $c = 0$ .

**Exercise 4.3. [Two independent Poissons]** Let  $X$  be a random variable taking values in  $\mathbb{N}_0$ , representing the number of users visiting server A within an hour. We assume that  $X \sim \text{Poisson}(\lambda)$  for some  $\lambda > 0$ .

- (a) Recall the definition of the Poisson distribution and argue why it is an appropriate probabilistic model for the described situation.
- (b) We know that the server crashes if at least 1000 people visit it within an hour. Express the probability of this event as a function of  $\lambda > 0$ .

- (c) Let  $Y$  be a random variable taking values in  $\mathbb{N}_0$ , representing the number of users visiting server B within an hour. Assume that  $Y \sim \text{Poisson}(\gamma)$  for some  $\gamma > 0$  and that  $X$  and  $Y$  are independent. That is,

$$\mathbb{P}[X = i, Y = k] = \mathbb{P}[X = i]\mathbb{P}[Y = k]^{(*)}, \quad i, k \in \mathbb{N}_0.$$

Find the probabilities  $\mathbb{P}[Z = k]$  for  $k \in \mathbb{N}_0$ , where  $Z := X + Y$ . What is the distribution of the random variable  $Z$ ?

**Hint:** It might be helpful to use the binomial formula

$$(a + b)^k = \sum_{i=0}^k \binom{k}{i} a^{k-i} b^i, \quad \text{for } a, b \in \mathbb{R}, k \in \mathbb{N}_0.$$

(\*) Similarly to the Remark following Definition 2.5 in the lecture notes, one can show that if  $X$  and  $Y$  take values in  $\mathbb{N}_0$ , then independence is equivalent to this formula. However, it is crucial that  $X$  and  $Y$  have **discrete distributions** (i.e., they are supported on a countable set), as this statement does not hold in general.

**Exercise 4.4. [On independence]** Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space and let  $A, B$  and  $A_i, i \in \{1, \dots, n\}$ , be events in  $\mathcal{F}$ . We assume that the  $A_i$  are pairwise disjoint.

- (a) Show that  $A$  and  $B$  are independent if and only if  $A$  and  $B^c$  are independent, which is also equivalent to the independence of  $A^c$  and  $B^c$ .
- (b) Show that if  $A$  and each  $A_i$  are pairwise independent for all  $i \in \{1, \dots, n\}$ , then  $A$  and  $\bigcup_{i=1}^n A_i$  are independent.
- (c) Assume that  $\mathbb{P}[A] = 1$ . Show that  $A$  and  $B$  are independent for all  $B \in \mathcal{F}$ .

**Remark:** Try to consider whether the results are intuitive and what their interpretation is.

**Exercise 4.5. [Studying pays off]** We have a representative student of the Probability Theory and Statistics course, whose diligence is represented by a random variable  $X_1$  taking values in  $\{0, 1\}$  with distribution  $\mathbb{P}[X_1 = 0] = 1 - \mathbb{P}[X_1 = 1] = 4/10$ . Here, we interpret  $\{X_1 = 1\}$  = “The student studies diligently” and  $\{X_1 = 0\}$  = “The student does not study diligently.”

Further, we define  $X_2$  taking values in  $\{0, 1\}$  as the random variable that represents whether the student attempts to pass the exam, where  $\{X_2 = 0\}$  = “The student does not register for the exam” and  $\{X_2 = 1\}$  = “The student registers for the exam.” We assume that

$$\mathbb{P}[X_2 = 1|X_1 = x_1] = 1 - \mathbb{P}[X_2 = 0|X_1 = x_1] = \frac{2 + 3x_1}{5}, \quad x_1 \in \{0, 1\}.$$

Finally, we define  $X_3$  taking values in  $\{0, 1\}$  as the random variable that represents whether the student actually passes the exam, where  $\{X_3 = 0\}$  = “The student does not pass the exam” and  $\{X_3 = 1\}$  = “The student passes the exam.” We assume that

$$\mathbb{P}[X_3 = 1|X_2 = x_2, X_1 = x_1] = \begin{cases} \frac{1+3x_1}{5}, & x_1 \in \{0, 1\}, x_2 = 1, \\ 0, & \text{otherwise.} \end{cases}$$

- (a) Describe in words the various conditional probabilities given in the problem, and specify their values.

- (b) At the end of the exam session, we know that the student did not pass the exam (i.e., either failed or did not register). What is the conditional probability that the student did not study diligently? That is, compute  $\mathbb{P}[X_1 = 0 | X_3 = 0]$ .

**Exercise 4.6. [Funny dice]** We have two different dice; on one of them, the 6 is replaced by a 7, and the other one is standard. First, we toss an unfair coin, which shows heads with probability  $p \in (0, 1)$ . If heads occurs, we choose the die with the 6; if tails occurs, we choose the die with the 7. Then, we roll the chosen die twice and compute the sum  $Y$  of the obtained face values.

- (a) Compute the probability for {the sum of the dice is 10} and {the sum of the dice is 12} as a function of  $p$ .
- (b) Determine the conditional probabilities of heads given that the sum is 10 and given that the sum 12.
- (c) Which of the events {coin toss results in heads}, {sum of the dice is 10}, and {sum of the dice is 12} are independent?