

## Solutions to problem set 1

1. (a) Consider the diagram

$$\begin{array}{ccc} H_n(\Delta^n, \partial\Delta^n) & \xrightarrow{\cong} & \tilde{H}_{n-1}(\partial\Delta^n) \\ \downarrow \cong & & \\ H_n(\Delta^n/\partial\Delta^n, *) & & \end{array} \quad (1)$$

The horizontal map is the boundary map from the (reduced) LES for the pair  $(\Delta^n, \partial\Delta^n)$ , which is an isomorphism by looking at the neighbouring terms in the LES. The vertical map is induced by the quotient map  $(\Delta^n, \partial\Delta^n) \rightarrow (\Delta^n/\partial\Delta^n, *)$  and is an isomorphism since  $(\Delta^n, \partial\Delta^n)$  is a good pair.

Consider now the tautological  $n$ -simplex  $\alpha_n : \Delta^n \rightarrow \Delta^n$ , which defines a class  $[\alpha_n] \in H_n(\Delta^n, \partial\Delta^n)$ . The image of  $[\alpha_n]$  under the vertical map is  $[\sigma_n] \in H_n(\Delta^n/\partial\Delta^n, *)$ , while its image under the horizontal map is the class  $[\beta_{n-1}] \in \tilde{H}_{n-1}(\partial\Delta^n)$  with

$$\beta_{n-1} = \partial_n \alpha_n = \sum_{i=0}^n (-1)^i F_i^n \in C_{n-1}(\partial\Delta^n),$$

where  $F_i^n : \Delta^{n-1} \rightarrow \partial\Delta^n$  is the  $i$ -th face map of the simplex  $\Delta^n$ . So once we know that  $[\beta_{n-1}]$  generates  $\tilde{H}_{n-1}(\partial\Delta^n)$ , we can conclude from (??) that  $[\sigma_n]$  generates  $H_n(\Delta^n/\partial\Delta^n, *)$ .

It is clear that  $[\beta_0]$  generates  $\tilde{H}_0(\partial\Delta^1)$ , so we know that  $[\sigma_1]$  generates  $H_1(\Delta^1/\partial\Delta^1, *)$ , which is what the problem asks us to prove for  $n = 1$ . We now proceed by induction; for the inductive step, consider the map  $\phi : \partial\Delta^n \rightarrow \Delta^{n-1}/\partial\Delta^{n-1}$  which collapses all except the zero-th face to a point, and the induced map  $\phi_* : H_{n-1}(\partial\Delta^n) \rightarrow H_{n-1}(\Delta^{n-1}/\partial\Delta^{n-1}, *)$ . Observe that  $\phi_*[\beta_{n-1}] = [\sigma_{n-1}]$ ; since  $[\sigma_{n-1}]$  generates by inductive assumption, we conclude that  $[\beta_{n-1}]$  generates.

- (b) Analogous to (a). In summary, there are isomorphisms

$$H_n(\Delta^n/\partial\Delta^n, *; G) \xrightarrow[\cong]{q_*} H_n(\Delta^n, \partial\Delta^n; G) \xrightarrow[\cong]{\partial_*} \tilde{H}_{n-1}(\partial\Delta^n; G) \xrightarrow[\cong]{\phi_*} H_{n-1}(\Delta^{n-1}/\partial\Delta^{n-1}, *; G)$$

$$[g\sigma_n] \longleftarrow [g\alpha_n] \longrightarrow [g\beta_{n-1}] \longrightarrow [g\sigma_{n-1}]$$

and

$$\begin{aligned} G &\longrightarrow \tilde{H}_0(\partial\Delta^0; G) \\ g &\longmapsto [g\beta_0]. \end{aligned}$$

2. Consider the cover of  $Y$  given by the subsets  $A = \Delta_+^n$  and  $B = \Delta_-^n$ . Both are contractible and we have  $A \cap B = \partial\Delta^n$ , so that the relevant piece of the corresponding reduced MV sequence reads

$$0 \rightarrow \tilde{H}_n(Y) \xrightarrow{\partial_*} \tilde{H}_{n-1}(\partial\Delta^n) \rightarrow 0$$

Note that  $\partial_*[\tau_+ - \tau_-] = [\partial\tau_+] = [\beta_{n-1}] \in \tilde{H}_{n-1}(\partial\Delta^n)$  with  $\beta_{n-1} \in C_{n-1}(\partial\Delta^n)$  defined as in the solution to the previous problem. Since  $[\beta_{n-1}]$  generates (see the previous problem) we deduce that  $[\tau_+ - \tau_-]$  generates.

We give an alternative inductive proof that  $[\beta_n]$  generates  $\tilde{H}_{n-1}(\partial\Delta^n)$  using the Mayer-Vietoris sequence. For  $n = 0$  the statement is clear. For the inductive step, consider the cover of  $\partial\Delta^{n+1}$  given by  $A := \text{im } F_0^{n+1}$  and  $B := \partial\Delta^{n+1} \setminus \text{int } A$  (the interiors don't cover all of  $\partial\Delta^{n+1}$ , but that can be repaired by taking small thickenings of  $A$  and  $B$ ). Since both  $A$  and  $B$  are contractible, the corresponding reduced MV sequence splits into pieces of the form

$$0 \rightarrow \tilde{H}_n(\partial\Delta^{n+1}) \xrightarrow{\cong} \tilde{H}_{n-1}(A \cap B) \rightarrow 0$$

Note that we can identify  $A \cap B = \partial A$  with  $\partial\Delta^n$  via  $F_0^{n+1}|_{\partial\Delta^n}$ . By definition of the MV boundary map  $\partial_* : \tilde{H}_n(\partial\Delta^{n+1}) \rightarrow \tilde{H}_{n-1}(A \cap B)$ , we have  $\partial_*[\beta_n] = [\partial F_0^{n+1}]$ , which in our identification  $A \cap B \cong \partial\Delta^n$  is  $[\beta_{n-1}]$ . Since  $\partial_*$  is an isomorphism and  $[\beta_{n-1}]$  generates  $\tilde{H}_{n-1}(\partial\Delta^n)$  by inductive assumption, it follows that  $[\beta_n]$  generates  $\tilde{H}_n(\partial\Delta^{n+1})$ .

3. (a) Let  $\tilde{\sigma} : \Delta^k \rightarrow X$  be a singular simplex. Then

$$T \circ \pi_c(\tilde{\sigma}) = \tilde{\sigma} + \Theta \circ \tilde{\sigma},$$

because  $\tilde{\sigma}$  and  $\Theta \circ \tilde{\sigma}$  are the two liftings of  $\pi \circ \tilde{\sigma}$ . Passing to homology, it follows that  $T_* \circ \pi_* = id + \Theta_*$ .

- (b) If  $H_i(X; \mathbb{Z}_2) \cong \mathbb{Z}_2$ , then  $\Theta_* : H_i(X; \mathbb{Z}_2) \rightarrow H_i(X; \mathbb{Z}_2)$  is the identity. ( $\Theta_*$  is an isomorphism and  $id$  is the only isomorphism on  $\mathbb{Z}_2$ .) So  $T_* \circ \pi_* = id + id = 0$  in degree  $i$ .
4. In the following, all homology groups have  $\mathbb{Z}_2$  coefficients. Given that  $H_k(\mathbb{R}P^n) = 0$  for  $k > n$  by assumption, the leftmost piece of the Smith sequence for the cover  $p : S^n \rightarrow \mathbb{R}P^n$  looks like

$$0 \rightarrow H_n(\mathbb{R}P^n) \xrightarrow{t_*} H_n(S^n) \xrightarrow{p_*} H_n(\mathbb{R}P^n) \xrightarrow{\partial_*} H_{n-1}(\mathbb{R}P^n) \rightarrow H_{n-1}(S^n) = 0 \rightarrow \dots$$

Here  $t_*$  is induced by the map  $C_*(\mathbb{R}P^n) \rightarrow C_*(S^n)$  taking a simplex  $\sigma : \Delta^k \rightarrow \mathbb{R}P^k$  to  $\tilde{\sigma} + \alpha \circ \tilde{\sigma}$ , where  $\tilde{\sigma} : \Delta^n \rightarrow S^n$  is one of the two possible lifts of  $\sigma$  to  $S^n$  and where  $\alpha : S^n \rightarrow S^n$  denotes the antipodal map. Note that we have  $t_* \circ p_* = (id + \alpha_*) : H_*(S^n) \rightarrow H_*(S^n)$ , which implies  $t_* \circ p_* = 0$  because  $\alpha_* = id : H_*(S^n) \rightarrow H_*(S^n)$  (because  $\alpha_*$  is an involution and  $H_k(S^n)$  either vanishes or is  $\mathbb{Z}_2$ ). This together with the fact that  $t_* : H_n(\mathbb{R}P^n) \rightarrow H_n(S^n)$  is injective implies that  $p_* : H_n(S^n) \rightarrow H_n(\mathbb{R}P^n)$  vanishes, and hence  $t_* : H_n(\mathbb{R}P^n) \rightarrow H_n(S^n) \cong \mathbb{Z}_2$  is an isomorphism. Moreover,  $p_* = 0$  implies that  $\partial_* : H_n(\mathbb{R}P^n) \rightarrow H_{n-1}(\mathbb{R}P^n)$  is an isomorphism, and the same is true for  $\partial : H_k(\mathbb{R}P^n) \rightarrow H_{k-1}(\mathbb{R}P^n)$  for  $k > 0$  since  $H_*(S^n) = 0$  except in degrees 0 and  $n$ . Inductively we obtain  $H_k(\mathbb{R}P^n) \cong \mathbb{Z}_2$  for all  $0 \leq k \leq n$ .

5. Recall that  $\pi_1(\mathbb{R}P^1) \cong \pi_1(S^1) \cong \mathbb{Z}$ ,  $\pi_1(\mathbb{R}P^n) \cong \mathbb{Z}_2$  and  $\pi_1(S^n) = 0$  for  $n > 1$ . Hence, if  $m = 1$  the only homomorphism  $\pi_1(\mathbb{R}P^n) \cong \mathbb{Z}_2 \rightarrow \pi_1(\mathbb{R}P^1) \cong \mathbb{Z}$  is the trivial homomorphism. So from now on we may assume that we have  $n > m > 1$ .

$$\begin{array}{ccc} S^n & \xrightarrow{\tilde{f}} & S^m \\ p^n \downarrow & & \downarrow p^m \\ \mathbb{R}P^n & \xrightarrow{f} & \mathbb{R}P^m \end{array}$$

For any  $n > m > 1$  we have

$$f_{\#} \circ p_{\#}^n(\pi_1(S^n)) = \{1\} = p_{\#}^m(\pi_1(S^m))$$


and  $f \circ p^n : S^k \rightarrow \mathbb{R}P^m$  always lifts to a map  $\tilde{f} : S^n \rightarrow S^m$ .

A generator of  $\pi_1(\mathbb{R}P^n)$  is represented by a loop that lifts to a path in  $S^n$  connecting two antipodal points (see also Hatcher example 1.43). The homomorphism  $f_\# : \pi_1(\mathbb{R}P^n) \cong \mathbb{Z}_2 \rightarrow \pi_1(\mathbb{R}P^m) \cong \mathbb{Z}_2$  can either be an isomorphism or trivial.

$$\begin{aligned}
& f \text{ induces an isomorphism } f_\# \\
& \iff \forall \text{ path } \gamma : [0, 1] \rightarrow S^n \text{ connecting antipodal points:} \\
& \quad f_\#([p^n \circ \gamma]) = [f \circ p^n \circ \gamma] = p_\#^m[\tilde{f} \circ \gamma] \in \pi_1(\mathbb{R}P^m) \setminus \{0\} \cong \mathbb{Z}_2 \setminus \{0\} \\
& \iff \forall \text{ path } \gamma : [0, 1] \rightarrow S^n \text{ connecting antipodal points:} \\
& \quad \tilde{f} \circ \gamma : [0, 1] \rightarrow S^m \text{ connects antipodal points} \\
& \iff \text{the lift } \tilde{f} : S^n \rightarrow S^m \text{ is equivariant.}
\end{aligned}$$

But, since  $n > m$ , by Bredon Theorem 20.1 the map  $\tilde{f}$  *cannot* be equivariant. Therefore, the induced map  $f_\#$  must be trivial.

6. Assume that  $r : \mathbb{R}P^3 \rightarrow \mathbb{R}P^2$  is a retraction and denote by  $i : \mathbb{R}P^2 \hookrightarrow \mathbb{R}P^3$  the inclusion. Then we have  $r \circ i = \text{id}_{\mathbb{R}P^2}$  and hence  $(r \circ i)_\# = \text{id} : \pi_1(\mathbb{R}P^2) \rightarrow \pi_1(\mathbb{R}P^2)$ , which is non-zero because  $\pi_1(\mathbb{R}P^2) = H_1(\mathbb{R}P^2; \mathbb{Z}) = \mathbb{Z}_2$ . On the other hand, we have  $(r \circ i)_\# = r_\# \circ i_\# = 0$  since  $r_\# = 0$  by the previous exercise. That is a contradiction.
7. Cf. the proof of Borsuk-Ulam in [Hatcher, pp. 174-176]!
8. Let  $n > m$  and supposed that there exists an equivariant map  $\phi : S^n \rightarrow S^m$ , i.e., such that  $\phi(-x) = -\phi(x)$  for all  $x$ . Consider the map  $f : S^{m+1} \rightarrow \mathbb{R}^{m+1}$  obtained by composing the restriction of  $\phi$  to  $S^{m+1} \subseteq S^n$  with the inclusion  $S^m \hookrightarrow \mathbb{R}^{m+1}$ . This map satisfies  $f(-x) = -f(x)$  for all  $x \in S^{m+1}$ . Since  $f(x) \in S^m$  and hence  $f(x) \neq -f(x)$ , we conclude  $f(-x) \neq f(x)$  for all  $x \in S^{m+1}$ , which contradicts the Borsuk-Ulam theorem.
9. Cf. [Bredon, Corollary IV.20.4]!
10. (a) For  $z \in \mathbb{R}P^k$  choose  $x \in B_+^k$  such that  $z = [x]$ . We define  $\phi(z)$  to be the point in  $S^k$  obtained from moving  $x$  down towards the South Pole  $S$  doubling the distance to the North Pole. (Explicitly for e.g.  $S^2$ , write  $x$  in spherical coordinates  $(\varphi, \theta)$  and define  $\phi(z) = (\varphi, 2\theta) \in S^k$ .)  $\phi : \mathbb{R}P^k \rightarrow S^k$  descends to a homeomorphism  $\mathbb{R}P^k / \mathbb{R}P^{k-1} \rightarrow S^n$ .



$f$  maps  $\text{Int}(B_\pm^k)$  homeomorphically onto  $S^k \setminus \{S\}$ .

- (b) The North Pole  $N$  has two preimages under  $f$ :  $N$  and  $S$ . Near  $N$ ,  $f$  is an orientation-preserving homeomorphism and hence the local degree at  $N$  is 1. Near  $S$ ,  $f$  is the composition of the antipodal map with  $f$  near  $N$ . Hence the local degree at  $S$  is  $(-1)^{k+1}$ . We conclude  $\deg(f) = 1 + (-1)^{k+1}$ .

*Remark.* There are many choices for  $\phi$ . One could for example also define  $\phi'$  using  $\phi'(z) = (-\varphi, 2\theta)$ . Then  $f$  has local degree  $-1$  near  $N$  and local degree  $-(-1)^{k+1}$  near  $S$ . So for that choice,  $\deg(f) = -(1 + (-1)^{k+1})$ .

However, for any choice of  $\phi$  as in (a), one has  $\deg(f) = \pm(1 + (-1)^{k+1})$ . The reason is, that  $f$  is a homeomorphism near  $N$  and a homeomorphism near  $S$ . Moreover, these

two homeomorphisms are related by the antipodal map because  $f$  factors through  $\mathbb{R}P^k$ . So if one of the local degrees is 1, then the other local degree will be  $(-1)^{k+1}$  and if one of the local degrees is  $-1$ , then the other local degree will be  $-(-1)^{k+1}$ .

- (c) We define a homeomorphism  $g: \mathbb{R}P^k \cup_{h_\partial} B^{k+1} \rightarrow \mathbb{R}P^k$ .  $g$  is the identity on  $\mathbb{R}P^k$ . To define what  $g$  does on  $B^{k+1}$ , let  $j: B^{k+1} \rightarrow S^{k+1}$  be the inclusion of  $B^{k+1} \approx B_+^{k+1}$  into  $S^{k+1}$ . Then  $g$  is defined to be  $q \circ j$  on  $B^{k+1}$ . One can check, that this gives a well-defined continuous bijective map  $g: \mathbb{R}P^k \cup_{h_\partial} B^{k+1} \rightarrow \mathbb{R}P^{k+1}$ . Hence  $h$  is a homeomorphism.
- (e) We compute

$$d(e^{(k+1)}) = \deg(f)e^{(k)} = (1 + (-1)^{k+1})e^{(k)} = \begin{cases} 2e^{(k)} & \text{if } k \text{ is odd} \\ 0 & \text{if } k \text{ is even.} \end{cases}$$